

SELFAT-53.5%

FUNGICIDE WETTABLE POWDER

Class :
Inorganic compound ⁽¹⁾
Dithiocarbamate ⁽²⁾

Foliar fungicide with contact, preventive action. Used to control many fungal diseases such as: Downy mildew, Blights, Leaf Spots and Scabs.

COMPOSITION : Each kg contains:

COPPER OXYCHLORIDE ⁽¹⁾	37.5% W/W (A.I.)
ZINEB ⁽²⁾	16% W/W (A.I.)

PROPERTIES:

SELFAT-53.5%: Foliar fungicide with contact, preventive action. Deposits must be on the crop before fungal spores begin to germinate .

SELFAT-53.5%: Used to control many fungal diseases such as: Downy mildew, Blights, Leaf Spot and Scab.

SELFAT-53.5%: Non-phytotoxic, except for Zinc-sensitive plants such as Tobacco and Cucurbits, and may cause injury to Plum and Peach foliage, especially at low temperatures.

USAGE AND DOSAGE:

CROP	DISEASES	APPLICATION RATE / 20 LT. WATER	SAFETY PERIOD
Fruit trees	Scab, Leaf spot.	60 – 80 gm	15 days
Potatoes	Late blight.	50 gm	7 days
Tomatoes	Late blight, Downy mildew.	40 – 60 gm	7 days

CAUTION: Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD: 48 hours after the last application.

COMPATIBILITY: Compatible with most common pesticides, but it is favorable to make compatibility test before use.

- For more details about first aid & precautions please refer to first aid & precautions index .