

VACOMIL-PLUS 50%

FUNGICIDE
WETTABLE POWDER

Class :
Acylalanine⁽¹⁾
Inorganic compound³

Effective systemic preventive and curative fungicide to control a wide range of fungal diseases such as (Blight, Downy mildew) on a wide range of crops and medical plants.

COMPOSITION : Each kg contains :

METALAXYL⁽¹⁾ 15% W/W (A.I)
COPPER OXYCHLORIDE⁽²⁾ 35% W/W (A.I)

PROPERTIES :

VACOMIL - PLUS 50% : Is a curative and preventive fungicide to control a wide range of fungal diseases on a wide range of crops.

VACOMIL - PLUS 50% : Is effective against a wide range of fungal diseases that affected a wide range of crops.

VACOMIL - PLUS 50% : Not phytotoxic for treated plants except some species of Apples.

VACOMIL - PLUS 50% : Controls a wide range of diseases such as: Downy mildew, Blights, Leaf spot, Scab.

VACOMIL - PLUS 50% : Is also used as seed treatment for number of crops.

USAGE AND DOSAGE :

| CROP | DISEASE | APPLICATION RATE GM/20 LT. WATER | SAFETY PERIOD (DAYS) |
|-----------------------------------|--------------------------------|-------------------------------------|----------------------------|
| Cauliflower, Cabbage. | Downy mildew. | 30 - 35 | 14 |
| Watermelon, Potato, Muskmelon. | Early and Late blight. | 30 - 35 | 14 |
| Grapes. | Downy mildew. | 30 - 40 | 12 |
| Medicinal plant. | Late blight , Downy mildew. | 30 | 20 |

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest, some varieties of red Apple are sensitive to this product. Sensitivity test should be done before application.

RE-ENTRY PERIOD : 48 hours after the last application.

COMPATIBILITY : Compatible with most other pesticides, but incompatible with Mercuric compounds, DNOC, Lime sulfur, and Dithiocarbamates.

* OTHER FORMULATION:

VACOMIL-PLUS - 76.5% WP
(METALAXYL 15% + COPPER OXYCHLORIDE 61.5% (Cu 35%)).

- For more details about first aid & precautions please refer to first aid & precautions index .