

# CURTINE-V

FUNGICIDE  
WETTABLE POWDER

Class :  
Dithiocarbamate <sup>(1)</sup>  
Cyanoacetamide oxime<sup>2)</sup>

Effective fungicide to control a broad spectrum of fungal diseases such as Downy mildew and Blights.

COMPOSITION : Each kg contains :

MANCOZEB <sup>(1)</sup>	46.5 %	W/W (A.I.)
CYMOXANIL <sup>(2)</sup>	4 %	W/W (A.I.)

#### PROPERTIES :

**CURTINE-V** : Highly absorbed by Plants, ensuring maximum utilization of the pesticide before being loosed by rain.

**CURTINE-V** : Protective Fungicide, prevent the formation of Mycelium at the incubation period and 4 days after infection.

**CURTINE-V** : Due to its dual structure its exhibits a tremendous efficacy against primitive fungi such as Downy mildew on Grapes, Lettuce, Onion, Tobacco, and Blights on Tomato and Potato.

**CURTINE-V** : Could be sprayed three days after rain to maintain good results.

#### USAGE AND DOSAGE :

CROP	DISEASE	APPLICATION RATE / 20 LT. WATER	SAFETY PERIOD (DAYS)
Grapes.	Downy mildew.	40-50 g	14
Potatoes And Tomatoes.	Late blight, Downy mildew.	40-50 g	15
Lettuce.	Downy mildew.	40-50 g	15
Onion, Garlic.	Downy mildew.	40-50 g	14
Tobacco.	Downy mildew.	40-50 g	14

**CAUTION:** Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD** : 24 hours after the last application.

#### COMPATIBILITY :

Compatible with most other pesticides, compatibility test is recommended before use.

#### \* OTHER FORMULATIONS:

CURTINE-V - 72% WP ( MANCOZEB 64% + CYMOXANIL 8% ),

CURTINE-V - 76.42 % WP ( MANCOZEB 70.5 % + CYMOXANIL 5.92% ).

- For more details about first aid & precautions please refer to first aid & precautions index .