

AMETRABOLE-80%

HERBICIDE
WETTABLE POWDER

Class :
1,3,5-triazine

A selective systemic herbicide used as pre and post -emergence to control most of annual grasses and broad-leaved weeds. Absorbed by the leaves and roots on a wide range of crops.

COMPOSITION : Each kg contains:

AMETRYN

80% W/W (A.I)

PROPERTIES:

AMETRABOLE-80%: A selective systemic herbicide used as pre and post-emergence to control annual grasses and broad-leaved weeds in Sugar cane field.

AMETRABOLE-80%: Controls many grasses in Sugar cane field, Citrus, Banana & Corn.

AMETRABOLE-80%: Compatible with most other pesticides, fertilizers at the normal application rates.

AMETRABOLE-80%: Does not prevent germination but kills weed after being absorbed to the root system.

USAGE AND DOSAGE:

CROP	DOSAGE	NOTES
Sugar cane	3 kg / ha + 500 ml (Spreading Agent)	- For control annual and broad leaved weeds apply to the soil surface as a pre or early post- emergence application.
	Mixture of 2 kg / ha AMETRABOLE-80% + 2.5 lt of 2,4-D (Amine)	- Integrated control against <i>Cyperus esculentus</i> and for late control, should be made before flowering.
Citrus	2 kg / ha	Apply as a direct spray in spring season. The second application after 60 - 90 days.
Bananas	2 kg / ha	Apply as a direct spray after fruit setting, repeat at 3-4 month intervals.
Corn	2.5 - 3 kg / ha	Apply as a direct spray, when the Corn at 18 cm height at least and never use before 6 weeks from planting.

Hectare = 10000 m²

CAUTION: Used only on above mentioned crops.

REMARKS:

- May be used in non-crop land to control most annual grasses and broad-leaved weeds.
- In order to achieve best results apply **AMETRABOLE-80%** at early post- emergent or when the plant at 5 cm height or 3 true leave stage.
- When used **AMETRABOLE-80%** as post-emergence its preferable to add spreading agent 0.1-0.3% of the solution to absorbed easily by leaves.
- **AMETRABOLE-80%** can be used as pre-emergence treatment, but soil must be humid to make easily movement of pesticide in weed root zone.
- Use 250-400 lt water /hectare when use back sprayer or traction sprayer, and use the high dosage rate for spraying weed when infestation is heavy.
- Good spray cover must be achieved to soil surface, its preferable to use nozzles specially for herbicide sprayer.

- For more details about first aid & precautions please refer to first aid & precautions index .

GRAZING RESTRICTIONS:

Don't graze or feed animals on the field which treated with pesticide newly.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most other pesticides, fertilizers at the normal application rates.

- For more details about first aid & precautions please refer to first aid & precautions index .