

TOVAP-30%

HERBICIDE
EMULSIFIABLE CONCENTRATE

Class :
Aryloxyphenoxypropionate⁽¹⁾
N.A.⁽²⁾

A selective systemic herbicide used to Control Wild oats, Canary grass and Annual Ryegrass in Wheat fields.

COMPOSITION : Each Liter Contains :

CLODINAFOP-PROPARGYL ⁽¹⁾	24 % W/V (A.I)
CLOQUINTOCET-MEXYL ⁽²⁾	6 % W/V (A.I)

PROPERTIES :

TOVAP-30%: Selective systemic herbicide should be applied only when weeds are actively growing.

TOVAP-30%: Controls Wild oats, Canary grass and Annual Ryegrass in Wheat fields.

USAGE AND DOSAGE :

CROP	WEEDS	APPLICATION RATE / HECTARE	TIME OF APPLICATION
Wheat 2 leaf to late jointing Zadocks.	Wild oats	65 ml+ 1lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 5 leaf stage
		85 ml + 1lt. of sticking materials / 100 lt. Water.	When weeds at 2 leaf to mid Tillering stage
	Canary grass	85 ml + 1lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 3 leaf stage
		125 ml + 1lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 5 leaf stage
		160 ml + 1lt. of sticking materials / 100 lt. Water.	When weeds at 2 leaf to early tillering stage
	Annual Ryegrass	160 ml + 1lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 4 leaf stage
210 ml + 1lt. of sticking materials / 100 lt. water.		When weeds at 2 leaf to early tillering stage	

Hectar = 10000 m².

NOTE:

- Calibration of sprayer is important before application.
- Complete coverage while spraying is recommended.
- Do not apply if rainfall is expected within 2 hours.
- Do not apply to crops which are under stress. For example, due to very dry, very wet, frosty or diseased conditions, nutrient deficiency or high insect pressure.

COMPATIBILITY:

- It is preferable to make mixing trial before spraying.
- When **TOVAP-30%** is applied with compatible herbicides always use a recommended sticking product.
- The following herbicides are known to be incompatible with **TOVAP-30%**: Diuron, Triasulfuron, (Bromoxynil + Dicamba + MCPA) and all 2,4-D formulations. Herbicides which are incompatible with **TOVAP-30%** must be applied separately. Apply these herbicides after 10 days from last treatment of **TOVAP-30%**.
- **TOVAP-30%** is compatible with **ULTRACIDIN-40%**.

- For more details about first aid & precautions please refer to first aid & precautions index .