

REPELL - 80%

**BIRDS REPELLENT
WETTABLE POWDER**

**Class :
Quinone-anthraquinone
xanthone**

Protection of Vegetables, Fruit trees, other crops, Cereals and Feed-stuffs sites from birds.

COMPOSITION : Each kg contains :

ANTHRAQUINONE

80% W/W (A.I)

PROPERTIES :

REPELL-80%: Prevents harmful birds like Crows, Blackbirds, Sparrows, Starlings to feed on seeds, emerging crops, fruits, vegetables, etc.

REPELL-80%: Makes seeds, buds and fruits unpalatable to the birds and reduces feeding damage to a minimum. However, the reaction on the different bird species greatly varies according to the season and available food supplies.

REPELL-80%: Displays good plant tolerance and does not impair the flavour of seeds, fruits or vegetables.

USAGE AND DOSAGE:

- **SEED TREATMENT** (Treat immediately before seeding).

• **APPLICATION RATE:**

60 gm /100 kg seed of Wheat, Rye and Barley.

90 gm /100 kg seed of Maize, Oats, Peas and Broad beans.

• **METHOD OF TREATMENT:**

To treat 100 kg seed, mix 60-90 gm **REPELL-80%** with 200-400 ml water. Evenly spray this mixture on the outspread seeds.

- **APPLICATION ON CROPS:**

1-2 kg /100 lt. water.

- - Spray at first sign of bird damage appears on vegetables, fruit ripening seed of Cereals, Maize, Grapes, etc.
- - Thoroughly spray all parts of the plants threatened by the birds.

• **SAFETY PERIOD:**

The minimum waiting period between final application and harvest is 7 days.

- To force out and repel away the nuisance birds, Crows, Geese from their sites between houses, grains and feed-stuff stores and around airports. It is used as follows:

- **METHOD OF APPLICATION:**

Mix (1-2) kg **REPELL-80%** with 100 lt water, first add the product to the 50 lt water mix well then add the remained amount of water with well agitation. Spray the infected areas. Repeat if necessary.

- **For more details about first aid & precautions please refer to first aid & precautions index .**