

AMOXAN

Water Soluble Powder

COMPOSITION: Each gm contains:

Amoxicillin trihydrate

200 mg

PROPERTIES:

- A beta-lactam antibiotic of the aminopenicillins.
- Safe, less toxic and rapidly absorbed from the gastrointestinal tract.
- Acts by impairing development of bacterial cell walls.
- Has a bactericidal action against a wide range of Gm+ve and Gm-ve bacteria, attacking chickens, pigeons, ducks, pre-ruminating animals, dogs and cats: Staph, Strept, Coryne, Listeria, Clostridia, Erysipelothrix, Coli, Salmonella, Pasteurella, Moraxella, Klebsiella, Proteus, Haemophilus, Shigella, Actinobacillus, Leptospira, Borrelia, Spirochaeta.

INDICATIONS:

- CRD, Colibacillosis, Salmonellosis, Fowl cholera, Bovine haemorrhagic septicaemia, Clostridial infections, Omphalitis and yolk sac infection, Coryza, Sinusitis, Air succulitis, Diarrhea, Enteritis, Spirochaetosis, Listeriosis, Erysipelas.
- Other respiratory, gastrointestinal and urogenital infections.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

DOSAGE:

(Firstly, dissolve well in 2 litres of water, then add to the water bulk to be medicated).

■ **Treatment:** (Duration: 5 days)

* **Poultry:** 100 gm / 200 litres of drinking water.

* **Lambs, Kids, Calves** (Pre-ruminating):

2 gm / 50 kg B. wt. twice daily and orally with drinking water.
(8 mg activity / kg B. wt.)

* **Dogs, Cats:**

0.5 gm / 10 kg B. wt. twice daily and orally with drinking water.
(10 mg activity / kg B. wt.)

* **Pigeons, Ducks:**

0.5 gm / 5 kg B. wt. with drinking water.
(20 mg activity / kg B. wt.)

■ **Prevention:** Half of the dosage for the same period.

SAFETY PERIOD:

- * For meat consumption: 1 week.
- * For eggs consumption: 3 weeks.

REMARKS:

- * Not used for table-eggs layers.
- * During treatment, eggs not used for human consumption.
- * Not used for rabbits.

STORAGE: Keep in dry, dark place at (15-30) °C.

PACKING: (100, 250, 500, 1000) GM.