

AMOXYCOL

Water Soluble Powder

COMPOSITION: Each gm contains:

Amoxicillin trihydrate	200	mg
Colistin sulphate	800,000	I.U

PROPERTIES:

- A combination of two wide - spectrum antibiotics which leads to a strong and synergistic action against poultry and livestock diseases.
- **AMOXYCOL** is used in a large-scale variety of infections of respiratory, alimentary, and urogenital tracts caused by Gm + ve and Gm – ve pathogenic bacteria such as: Coli, Salmonella, Clostridia, Mycoplasma, Pasteurella, Haemophilus, Proteus, Actinomyces, Staphylococcus, Streptococcus, Coryne, Listeria, Yersinia, Moraxella, Bordetella, Actinobacillus, Klebsiella etc....

INDICATIONS:

- Salmonellosis, Colibacillosis, Clostridial infections, CRD, Omphalitis and Yolk sac infection, Sinusitis, Coryza, Air Sacculitis, Pasteurellosis, Mycoplasmosis, Leptospirosis.
- Cases of enteritis, septicaemia, secondary bacterial and generalized infections.
- Respiratory, alimentary and uro-genital infections in other animals.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

DOSAGE:

(Firstly, dissolve well in 2 litres of water, then add to the water bulk to be medicated).

- **Treatment:** (Duration: 3 - 5 days)
 - * **Poultry:** 100 gm per 200 litres of drinking water.
 - * **Horses:** 8-10 gm per animal twice daily and orally with water.
 - * **Pre-ruminating calves, Foals:** 3-6 gm per animal twice daily and orally with water.
 - * **Lambs, Kids :** 1-3 gm per animal twice daily and orally with water.
- **Prevention:** Half dosage for the same period.

SAFETY PERIOD:

- * For meat consumption: one week.
- * For eggs consumption: 3 weeks.

REMARKS:

- * Not used for table-eggs layers.
- * During treatment eggs not used for human consumption.
- * Not used for rabbits.

STORAGE: Keep in dry, dark place at (15-30) °C.

PACKING: (100, 250, 500, 1000) GM.