

AMPIVAP AMPIVAP-50%

Water Soluble Powder
For Poultry, Game birds, and other animals

COMPOSITION: Each gm contains:

	Ampivap	Ampivap-50%
Ampicillin trihydrate	300 mg	500

PROPERTIES: Ampicillin is a very active antibiotic against a large scale of Gm+ ve and Gm- va bacteria such as: Coli, Salmonella, Staph, Strept, Clostridia, Pasteurella, Chlamydia, Mycoplasma, Spirochaeta, Coryne, Haemophilus gallinarum etc....

INDICATIONS:

■ **Poultry (Chickens, Turkeys), Game birds:**

- * Colibacillosis, CRD, Sinusitis, Necrotic and Ulcerative enteritis (Clostridia).
Omphalitis and Yolk sac infection, Coryza, Salmonellosis, Spirochaetosis, Fowl cholera (Pasteurellosis) Staphylococcal arthritis.
- * Stress Conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.
- * Recommended as supportive treatment against secondary bacterial complications in cases of viral infections.

■ **Lambs, Calves, Foals:** Bacterial enteritis, scour, infections of respiratory and uro-genital tracts, preventive against mortality of newly-born and young animals, Haemorrhagic septicaemia (Pasteurella).

■ **Dogs, Cats:** Bacterial gastro-enteritis, Septicaemia, Infections of respiratory and uro-genital tracts, Joint ill, Leptospirosis.

DOSAGE:

(Firstly, dissolve well in 2 litres of water, then add to the water bulk to be medicated).

■ **Treatment:** Duration for 3-5 days

- * **Chickens, Turkeys, Game birds** (Quails, Partridges, Pheasants, Guinea fowls).
- * **Ampivap:** 100 gm per 200 litres of drinking water.
- * **Ampivap-50%:** 60 gm / 200 litres.
- * **Other animals:**
 - **Ampivap:** 0.35-0.70 gm per 10 kg B.wt. twice daily and orally with water.
 - **Ampivap-50%:** 0.2-0.4 gm / 10 kg B.wt. twice daily and orally with water.

■ **Prevention:** Half dosage for the same period.

SAFETY PERIOD:

- * For meat consumption: 15 days.
- * For eggs consumption: 3 weeks.

REMARKS:

- * Not used for rabbits.
- * Not used for table-eggs layers.
- * During treatment, eggs not used for human consumption.

STORAGE: Keep in dry, dark place at (15-30) °C.

PACKING: (100, 250, 500, 1000) GM.