

ENROSOL-S P/FORTE/20%

Powder and Solution

COMPOSITION: Each (gm,ml) contains:

	<u>S/P</u>	<u>Forte (Sol)</u>	<u>20 % (Sol)</u>
Enrofloxacin	100 mg	120 mg	200 mg

PROPERTIES:

- The effective anti-bacterial agent which belongs to fluoroquinolone group.
- Characterised by quick absorption and rapid high serum level.
- Has a broad spectrum efficacy against a wide range of bacteria and mycoplasmas.

INDICATIONS:

- **Poultry:** CRD, Colibacillosis, Coryza, Fowl cholera, Fowl typhoid, other Salmonellosis (S.typhimurium, S.enteritidis etc.), Staph and Strept infections, Mycoplasmosis, Omphalitis and yolk Sac infection, and Erysipelothrix infections of tyrkeys.
- **Calves, Lambs:** Respiratory tract infections (Pasteurella, Mycoplasma), enteritis, dysentery, diarrhea of newly-born and growing off-springs, caused by (Colibacillosis, Salmonellosis), and Haemophilus.
- **Rabbits:** Pasteurellosis, Enterotoxaemia, Enteritis, dysentery, respiratory tract infections.
- **Piglets:** Respiratory tract infections, diarrhea and enteritis of newly-borns, caused by bacteria (E. Coli, Salmonella etc.)..

DOSAGE:

■ Treatment:

* Poultry:

- **Enrosol S/P:** 100 (gm,ml) per 200 litres of drinking water for 3 days.
- **Enrosol Forte:** 85-100 ml/200 litres of drinking water for 3 days.
- **Enrosol 20% :** 50 ml/ 200 litres of drinking water for 3 days.
- In cases of salmonella, continue treatment for additional 2 days.

* Calves, Lambs, Kids, Rabbits, Piglets: (3-5 mg activity / kg B.wt.).

- **Enrosol S/P:** 0.3-0.5 (gm,ml) per 10 kg B.wt. daily and orally for 3-5 days.
- **Enrosol Forte:** 0.6-1 ml/25kg B.wt. daily and orally for 3-5 days.
- **Enrosol 20% :** 0.15-0.25 ml/10 kg B.wt. daily and orally for 3-5 days.

- **Prevention:** Half dosage for the same period.

SAFETY PERIOD: For meat consumption: 7 days.

STORAGE: Keep in a dry, dark place at (15-30) °C.

PACKING: (100, 250, 500, 1000) GM, ML.