

# CYPERMETHRIN-10 EC

## Pyrethroidal Insecticide Emulsifiable Concentrate

**COMPOSITION :** Each litre contains :

Cypermethrin

100 gm

**PROPERTIES :**

- Non-systemic insecticide with contact and stomach action.
- It is formulated for use as dip or spray.
- Well tolerated by farm animals.
- Biodegradable and does not accumulate in the tissues or in milk.
- Has rapid knockdown effects on all ticks including those resistant to organo-phosphorus and organo-chlorine insecticides.
- Very active at low concentration and has low mammalian toxicity.

**USES:**

- For the treatment of scab and mange mites of livestock beside poultry mites as well.
- For the control of different ectoparasites (ticks, fleas, lice, keds and screw worms) attacking cattle, sheep, goats and camels.
- For the control of nuisance flies and other insects in farms, stables and animals houses.

**METHODS OF APPLICATION:**By spraying or dipping after dilution with water as follows;

Species	Dilution Rate	<b>Treatment may be repeated after 1-2 weeks</b>
Cattle, Camels	1:1000	
Flies on cattle	1:300	
Sheep, goats	1:800	
Poultry	1:750 spray (60 litres per 1000 birds)	
Animals Houses	1:50	

**PRECAUTIONS:**

- Toxic to fishes and bees.
- Avoid application on sick, weak, exhausted and young animals.
- Avoid contact with eyes, mouth, skin and inhalation.
- Before spraying, wear protective clothes, boots, gloves and goggles.
- Do not eat, smoke or drink during spraying.
- Avoid polluting ponds, waterways and water sources.
- Don't allow off-spring to suckle until mothers are completely dried from the insecticide liquid.

**VAPCO**

**INSECTICIDES  
EMULSIFIABLE  
CONCENTRATES  
(EC)**

**FIRST - AID :**

■ **Eye contact :** Wash with water for (15) minutes.

■ **Skin contact:** Wash with soap and water.

■ **Inhalation:**

\* Move patient to fresh air.

\* Apply artificial respiration, give oxygen when necessary.

■ **Ingestion:**

\* Do not induce vomiting.

\* Apply gastric lavage.

■ Get medical help, when necessary.

**ANTIDOTE:**

\* Not specific, treat according to symptoms.

\* Give diazepam in case of convulsions.

**SAFETY PERIOD:**For meat and milk consumption: One week.

**STORAGE:**

■ Keep away from children, food and drinks.

■ Keep in a dry, dark place at (15-25) °C.

**PACKING:** (1,5, 20) Litres.