

# VAP MALATHION-50% EC

## Insecticide and Acaricide Emulsifiable Concentrate

**COMPOSITION:** Each litre contains:

Malathion

500 gm

**PROPERTIES:**

- Non-systemic insecticide and acaricide with contact, stomach and respiratory action.
- Strong organo-phosphorus ectoparasiticide which is effective at low concentration.
- Low mammalian toxicity.

**USES:**

- Treatment of different types of mange.
- Control of many species of ticks, mites, lice, keds, fleas, attacking livestock.
- Control of wound-infesting maggots which attack the animal.
- For the combat of insects in farms, stables and other animals houses.

**METHODS OF APPLICATION:**

Species	Dilution Rate	Remarks
Cattle	1:1100	Spray
Sheep, Goats	1:1100	Spray and dips
Equine	1:1100	Spray
Dogs	1:4000	Spray
Poultry	1:2100	Spray
Animals Houses	1 : 100	

**PRECAUTIONS:**

- Toxic to bees and fishes.
- Avoid contact with eyes, mouth, skin and inhalation.
- Wear protective clothes, gloves, boots and face shield before spraying.
- Do not eat, drink or smoke during spraying.
- Do not treat sick, exhausted, weak or very young animals.
- Avoid polluting water-sources, ponds and water canals.
- Don't allow off-spring to suckle until mothers are completely dried from the insecticide liquid.

**VAPCO**

**INSECTICIDES  
EMULSIFIABLE  
CONCENTRATES  
(EC)**

**FIRST - AID**

- **Eye contact:** Wash with water for (15) minutes.
- **Skin contact:** Wash with soap and water.
- **Inhalation:**
  - \* Move patient to fresh air.
  - \* Apply artificial respiration, give oxygen when necessary.
- **Ingestion:**
  - \* Induce vomiting by throat fingure touch only.
  - \* Apply gastric lavage.
- Get medical help, when necessary.

**ANTIDOTE:**

- \* Atropine, PAM, Toxogonin.
- \* Continue symptomatic treatment.

**SAFETY PERIOD:** For milk consumption: One day.

**STORAGE:**

- Keep away from children, food and drinks.
- Keep in a dry, dark place at (15-25) °C.

**PACKING:** (250, 1000 ) ML.  
(5 , 20) Litres.