

AMOXYCOL

Water Soluble Powder

A combination of two wide – spectrum antibiotics which leads to a strong and synergistic action against poultry and livestock diseases.

COMPOSITION: Each gm contains:

Amoxycillin trihydrate	200	mg
Colistin sulphate	800,000	I.U.

PROPERTIES:

AMOXYCOL is used in a large-scale variety of infections of respiratory, alimentary, and urogenital tracts caused by Gm + ve and Gm – ve pathogenic bacteria such as: Coli, Salmonella Clostridia, Mycoplasma, Pasteurella, Haemophilus, Proteus, Actinomyces, Staphylococcus, Streptococcus, Coryne, Listeria, Yersinia, Moraxella, Bordetella, Actinobacillus, Klebsiella etc....

INDICATIONS:

- Salmonellosis, Colibacillosis, Clostridial infections, CRD, Omphalitis and Yolk sac infection, Sinusitis, Coryza, Air Succulitis, Pasteurellosis, Mycoplasmosis, Leptospirosis.
- Cases of enteritis, septicaemia, secondary bacterial and generalized infections.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

DOSAGE: (Firstly, dissolve well in 2 litres of water, then add to the water bulk to be medicated).

■ **Treatment:**

- **Poultry:** 100 gm per 200 litres of drinking water for 3-5 days.
- **Cattle, Horses:** 8-10 gm per animal twice daily and orally with water.
- **Calves, Foals:** 3-6 gm per animal twice daily and orally with water.
- **Sheep, Goats:** 2-5 gm per animal twice daily and orally with water.

- **Prevention:** Half dosage for the same period.

SAFETY PERIOD:

- For meat consumption: 1 week.
- For milk consumption: 2 days.

■ **REMARKS:**

- Not used for layers.
- Milk taken during treatment is not used for human consumption.
- Not used for rabbits.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.