

AMPIVAP-30%

Water Soluble Powder

For Poultry, Game birds, Ruminants, Equines, Canines

COMPOSITION: Each gm contains:

Ampicillin Trihydrate 300 mg

PROPERTIES: Ampicillin is a very active antibiotic against a large scale of Gm+ ve and Gm- va bacteria such as: Coli, Salmonella, Staph, Strept, Clostridia, Pasteurella, Chlamydia, Mycoplasma, Spirochaeta, Coryne, Haemophilus, etc....

INDICATIONS:

- **Poultry (Chickens, Turkeys), Game birds:**
 - Colibacillosis, CRD, Sinusitis, Necrotic and Ulcerative enteritis (Clostridia).
Omphalitis and Yolk sac infection, Coryza, Salmonellosis, Spirochaetosis, Fowl cholera (Pasteurellosis) Staphylococcal arthritis.
 - Stress Conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.
 - Recommended as supportive treatment against secondary bacterial complications in cases of viral infections.
- **Ruminants, Equines:** Bacterial enteritis, scour in lambs, calves and foals, infections of respiratory and uro-genital tracts, preventive against mortality of newly-born and young animals, Haemorrhagic septicaemia (Pasteurella) in cattle.
- **Dogs, Cats:** Bacterial gastro-enteritis, Septicaemia, Infections of respiratory and uro-genital tracts, Joint ill, Leptospirosis.

DOSAGE: (firstly, dissolve well in 2 litres of water, then add to the water bulk to be medicated)

- **Treatment:** Duration for 3-5 days.
 - **Chickens, Turkeys, Game birds** (Quails, Partridges, Pheasants, Guinea fowls): 100 gm per 200 litres of drinking water.
 - **Other animals:**
 - 0.35-0.70 gm per 10 kg B.wt. twice daily and orally with water:
- **Prevention:** Half dosage for the same period.

SAFETY PERIOD: For meat consumption: 15 days.
For milk consumption: 3 days.

REMARKS:

- * Not used for rabbits.
- * Not used for table – eggs layers.
- * During treatment, eggs not used for human consumption.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.