

# **AMPIVAP-50%**

**Water Soluble Powder**

**For Poultry, Game birds, Ruminants, Equines, Canines**

**COMPOSITION:** Each gm contains:

Ampicillin Trihydrate

500 mg

**PROPERTIES:** Ampicillin is a very active antibiotic against a large scale of Gm+ ve and Gm- va bacteria such as: Coli, Salmonella, Staph, Strept, Clostridia, Pasteurella, Chlamydia, Mycoplasma, Spirochaeta, Coryne, Haemophilus, etc....

## **INDICATIONS:**

### ■ **Poultry (Chickens, Turkeys), Game birds:**

- Colibacillosis, CRD, Sinusitis, Necrotic and Ulcerative enteritis (Clostridia).  
Omphalitis and Yolk sac infection, Coryza, Salmonellosis, Spirochaetosis, Fowl cholera (Pasteurellosis) Staphylococcal arthritis.
- Stress Conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.
- Recommended as supportive treatment against secondary bacterial complications in cases of viral infections.

### ■ **Ruminants, Equines:** Bacterial enteritis, scour in lambs, calves and foals, infections of respiratory and uro-genital tracts, preventive against mortality of newly-born and young animals, Haemorrhagic septicaemia (Pasteurella) in cattle.

### ■ **Dogs, Cats:** Bacterial gastro-enteritis, Septicaemia, Infections of respiratory and uro-genital tracts, Joint ill, Leptospirosis.

**DOSAGE:** (firstly, dissolve well in 2 litres of water, then add to the water bulk to be medicated)

### ■ **Treatment:** Duration for 3-5 days.

- **Chickens, Turkeys, Game birds** (Quails, Partridges, Pheasants, Guinea fowls): 60 gm / 200 litres.
- **Other animals:** 0.2-0.4 gm / 10 kg B.wt. twice daily and orally with water.

### ■ **Prevention:** Half dosage for the same period.

**SAFETY PERIOD:** For meat consumption: 15 days.

For milk consumption: 3 days.

## **REMARKS:**

\* Not used for rabbits.

\* Not used for table – eggs layers.

\* During treatment, eggs not used for human consumption.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** (100, 250, 500, 1000) GM.