

Water Soluble Powder

The safe and effective anthelmintic against intestinal round worms

COMPOSITION: Each gm contains:

Piperazine citrate 360 mg

PROPERTIES:

- Has a wide margin of safety, it can be used for young or pregnant animals.
- Excellent efficacy against different species of intestinal nematodes (round worms) in poultry, ruminants, equines, rabbits, dogs and cats.

INDICATIONS:

Treatment and control of ascaridiosis due to intestinal nematodes (round worms) sush as: Ascaridia, Capillaria, Oxyuris, Strongylides, Oesophagostomum etc..., in poultry and other domestic animals.

DOSAGE:

• **Poultry:** (Treatment duration for 2 successive days).

(0.8-1) gm per bird daily.

(32 mg activity / kg body wt.).

(8-10 sachets of 100 gm per 1000 birds daily.

OR (0.8-1) kg per 1000 birds daily.

■ Sheep, Goats, Cattle, Camels, Equines:

0.8 gm per kg body wt.

(110 mg activity / kg body wt.).

■ Rabbits:

0.5-0.8 gm per kg body wt.

(70-105 mg activity / kg body wt.).

Dogs, Cats:

0.35-0.5 gm per kg body wt.

(45-65 mg activity / kg body wt.).

Repeat treatment after 4-6 weeks for one day.

SAFETY PERIOD: For meat consumption: 2 days.

STORAGE: Keep in a dry, dark place at below 30 °C.

PACKING: (100, 250, 500, 1000) ML.