

Water soluble and Feed Additive Powder Vitamins & Aminoacids for Poultry and Livestock

.

COMPOSITION: Each Kg contains:

VITAMINS			AMINO
Vitamin A	15,000,000	I.U.	Aspartic a
Vitamin D3	3,000,000	I.U.	Threonine
Vitamin E	20,000	I.U.	Serine
Vitamin K3	2	gm	Glutamic
Vitamin B1	2	gm	Proline
Vitamin B2	8	gm	Glycine
Vitamin B6	5	gm	Alanine
Vitamin B12	0.01	gm	Cystine
Calcium – D- pantothenate	12	gm	Valine
Nicotinic acid	40	gm	Methionin
Vitamin C	60	gm	Isoleucine
Folic acid	1	gm	Leucine
Biotin	0.6	gm	Tyrosine
			Phenylala
			Lysine
			Histidine

	AMINOACIDS		
	Aspartic acid	5720	mg
	Threonine	1540	mg
	Serine	2760	mg
ı	Glutamic acid	10,380	mg
ı	Proline	12,100	mg
ı	Glycine	23,850	mg
ı	Alanine	9120	mg
ı	Cystine	60	mg
ı	Valine	2500	mg
1	Methionine	800	mg
ı	Isoleucine	1220	mg
ı	Leucine	2970	mg
1	Tyrosine	520	mg
	Phenylalanine	2490	mg
	Lysine	3820	mg
	Histidine	560	mg
	Arginine	7310	mg
	Tryptophan	1200	mg
	Hydroxyproline	9580	mg

INDICATIONS:

- Hypovitaminosis.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.
- Maintains good performance and increases the production.
- Improvement of feed conversion, productivity, the growth, fertility and hatchability rates, feathering and flesh development.
- Enhancement of better rearing results.
- Helps in producing high eggs-quality, and bright, strong egg-shells.
- Increase of viability and resistance against diseases.
- Effective in cases of caged-layers fatigue.

DOSAGE: Duration for (3-5) days.

- **Poultry:** 100 gm per 200 litres of drinking water.
- Calves, Foals, Weaners: 1-2 gm daily and orally with water.
- Via the feed: 1 kg per ton.

REMARK:

To ensure complete dissolution, dissolve 100 gm (product) in at least 10 litres of water and then add to the water bulk.

STORAGE: Keep in a dry, dark place at (15-25) °C. **PACKING:** (100, 250, 500, 1000) GM.