

CIPROTRIL-10%

Oral Solution

COMPOSITION: Each ml contains:

Ciprofloxacin 100 mg

PROPERTIES:

- The effective anti-bacterial agent which belongs to fluroquinolone group.
- Characterised by quick absorption and rapid high serum level.
- Has a broad spectrum efficacy against a wide range of bacteria and mycoplasmas.

INDICATIONS:

- **Poultry:** CRD, Coli septicemia, Coryza, Fowl cholera, Sinusitis, Air succulitis, Fowl typhoid, other salmonellosis (*S. typhimurium*, *S. enteritidis* etc...), Staph and Strept infections, Omphalitis and yolk sac infection, Mycoplasmosis, and Erysipelothrix infection of turkeys.
- **Calves, Lambs:** Respiratory tract infections (*Pasteurella*, *Mycoplasma*), enteritis, dysentery, diarrhea of newly-born and growing off-springs, caused by (*Colibacillosis*, *Salmonellosis*), and *Haemophilus*, besides, uro-genital tract infections.
- **Rabbits:** Pasteurellosis, enterotoxaemia, enteritis, dysentery, respiratory tract infections.
- **Piglets:** Uro-genital and respiratory tract infections, diarrhea, and enteritis of newly-borns, caused by bacteria such as: *Coli*, *Salmonella* etc...

DOSAGE:

- **Treatment:**(Duration for 3-5 days).
 - **Poultry:** 100 ml per 200 litres of drinking water .
 - **Calves, Lambs, Rabbits, Piglets:** 0.3-0.5 ml / 10 kg B.wt. daily and orally with water. (3-5 mg active substance per kg B.wt. daily).
- **Prevention:** Half dosage for the same period.

SAFETY PERIOD: For meat consumption: 7 days.

REMARK: During treatment, don't medicate water with mineral preparations.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) ML.