

COCCIDOLE

Feed Additive Powder

COMPOSITION: Each gm contains:
Clopidol **125 mg**

PROPERTIES:

- Clopidol is of pyridinol group which acts on first generation schizonte coccidia, and thus prevents the disease occurrence.
- It does not develop any immunity against coccidiosis. It is non-toxic for non-host species, very safe to be used and is compatible with other medicaments including antibiotics, so it can be administered with other additives except other anticoccidials.
- It helps and improves metabolism, so it promotes growth and improves performance, feed conversion and weight gain.

INDICATIONS:

For prevention and treatment of most species of *Eimeria* strains causing caecal and intestinal coccidiosis in poultry, rabbits, turkeys and game birds (partridges, pheasants, guinea-fowls).

DOSAGE:

- **Prevention:** Start medication at day old of the flock.
- **Poultry:** 1 kg per ton of feed continuously up to 16 weeks old.
- **Rabbits:** 1.6 kg per ton of feed continuously.
- **Turkeys:** 800 gm per ton of feed continuously up to 12 weeks old.
- **Game birds:** 1 kg per ton of feed continuously.

- **Treatment:** According to the severity of the case, double the preventive dose up to 2-4 times for 2-4 days, then back to the preventive dose.

REMARK:

- Not recommended for layers.
- Not mixed with other anticoccidials.

SAFETY PERIOD: For meat consumption : 7 days.

STORAGE: Keep in a dry, dark place at below 30 °C.

PACKING: (1, 5, 10, 25) KG.