

# **CYCO – S**

## **Water Soluble and Feed Additive Powder**

**COMPOSITION:** Each gm contains:

N 1( 4,6 dimethylpyrimidine – 2 pyrazinyl ) sulphamide Na	320 mg
5 – Veratrylpyrimidine – 2,4 diamine	10 mg

**PROPERTIES:** An effective anticoccidial and antibacterial combination.

### **INDICATIONS:**

Prevention and treatment of Salmonella and most species of Eimeria strains causing caecal, hepatic and intestinal coccidiosis in poultry ( chickens, turkeys ) and rabbits.

### **DOSAGE:**

- **Poultry:** 100 gm per 200 litres of drinking water for 3 consecutive days, followed by plain water for 2 days, then apply medicated water for the next 2 days.
- **Rabbits:** 1 gm per 4 litres of drinking water for 3-5 days.
- **Via feed:** 2 kg per ton of feed for 3 days.
- **In severe cases:** The dose can be doubled and for the same period.

**SAFETY PERIOD:** For meat consumption: 7 days.

### **REMARKS:**

- Not recommended for laying flocks.
- Ensure adequate water intake during treatment to reduce the risk of crystalluria.

**STORAGE:** Keep in a dry, dark place at below 30 °C

**PACKING:** (100, 250, 500, 1000) GM.