

# ***CYCO – V***

---

## **Water Soluble Powder**

**COMPOSITION:** Each gm contains:

<b>Sulphaquinoxaline Sodium</b>	<b>160 mg</b>
<b>Diaveridine HCl</b>	<b>20 mg</b>

### **PROPERTIES:**

#### ■ **Sulphaquinoxaline (SQS):**

- Has anticoccidial effect and antibacterial activity against Salmonella, Pasteurella, E. coli, etc...It acts also against the second generation schizonts of coccidia.
- It is rapidly, completely absorbed and is slowly excreted from the body.
- Due to the slow elimination rate, it has a persistent action, which enhances its effect against coccidia.

#### ■ **Diveridine:**

- It is a coccidiostatic agent, and has also a bacteriostatic action.
- It forms with (SQS) a synergistic action against the different coccidia strains.

### **INDICATIONS:**

- For prevention and treatment of caecal and intestinal coccidia, bacterial enteritis and diarrhea in poultry ( chickens, turkeys ), rabbits and small ruminants.
- It is also used as an aid in controlling mortality in acute fowl cholera and fowl typhoid of poultry.

### **DOSAGE:**

#### ■ **Treatment:**

- **Poultry:** 100 gm per 200 litres of drinking water for 3 days, followed by plain water for 2 days, then apply medicated water for the next 3 days.
- **Rabbits:** 0.5 gm per litre of drinking water for 3-4 days.
- **Sheep, goats, calves:** 3 gm per 10 kg B. wt. for 3-4 days.

- **Prevention:** Half dosage for the same period.

**SAFETY PERIOD:** For meat consumption: 7 days.

### **REMARKS:**

- Not recommended for laying flocks and milking animals.
- Ensure adequate water intake during treatment to reduce the risk of crystalluria.
- Not given in cases of Gumboro infection, severe hepatic or renal dysfunctions.

**STORAGE:** Keep in a dry, dark place at below 30 °C.

**PACKING:** (100, 250, 500, 1000) GM.