

DIACOX

Water Soluble Powder

COMPOSITION: Each gm contains:

Sulphadimidine Na	800 mg
Diaveridine HCl	80 mg

PROPERTIES:

- A highly active and synergistic combination of anticoccidial and antibacterial agents.
- Effective against different *Eimeria* spp. (*Acervulina*, *Necatrix*, *Tenella*, *Brunetti*, *Maxima*, *Meleagrimitis*, *Adenoides*).
- It has no adverse effects on egg production, fertility and growth.

INDICATIONS:

For the treatment of salmonellosis, caecal, hepatic and intestinal coccidiosis in poultry, turkeys, guinea fowls and ducks.

DOSAGE:

■ **Poultry, Turkeys, Guinea fowls:**

50 gm per 200 litres of drinking water for 5 consecutive days or treat for 3 consecutive days, followed by plain water for two days, then medicate for the next 3 days.

- #### ■ **Ducks:**
- 100 gm per 200 litres of drinking water for 5 consecutive days or treat for 3 consecutive days followed by plain water for two days, then medicate for the next 3 days.

■ **SAFETY PERIOD:**

- For meat consumption: 14 days.
- For eggs consumption: 12 days.

REMARK:

Ensure adequate water intake during treatment to reduce the risk of crystalluria.

STORAGE: Keep in a dry, dark place at below 30 °C.

PACKING: (100, 250, 500, 1000) GM.