

# ***DIAZINON***

**(15 EC, 20 EC, 25 EC, 60 EC)**

**Organophosphorus Insciticide and Acaricide  
Emulsifiable Concentrate**

**COMPOSITION :** Each litre contains :

Diazinon 150 gm, 200 gm, 250 gm, 600 gm

## **PROPERTIES :**

- Non-systemic insecticide and acaricide with contact, stomach and respiratory action.
- A powerful and broad spectrum insecticide and acaricide.
- Well tolerated by farm animals and pets except cats and poultry.

## **USES:**

- Treatment of scab, mange and for the control of mites, ticks, lice, keds, fleas and screw worms on sheep, goats, cattle and camels.
- Control of stable flies and blow-fly larvae.

**METHODS OF APPLICATIONS:** By spraying or dipping after dilution with water as follows:

<b>Species</b>	<b>Spray</b>			
Sheep, goats, Cattle, Camels	<b>15 EC</b> 1:250	<b>20 EC</b> 1:350	<b>25 EC</b> 1:400	<b>60 EC</b> 1:1000
Animals houses	1:6	1:8	1:10	1:25
<b>DIP</b>				
Sheep, goats	1:600 Replenish 1:200	1:800 Replenish 1:250	1:1000 Replenish 1:350	1:2400 Replenish 1:800
Cattle, camels	1:250 Replenish 1:100	1:350 Replenish 1:150	1:400 Replenish 1:160	1:1000 Replenish 1:400

## **PRECAUTIONS:**

- Diazinon must be used directly after being diluted with water, and not kept for the next day to be used especially for cattle.
- In presence of water, the stored diazinon is oxidized to diazoxon which is harmful to animals especially cattle.
- Opened bottles should be used within 3 months.
- Do not use after expiration date.
- Toxic to fishes, bees, cats, poultry and birds.
- Avoid application on sick, exhausted, weak and young animals.
- Before spraying, wear protective clothes, boots, gloves and goggles.
- Do not eat, drink or smoke during spraying.
- Avoid polluting ponds, waterways and water sources.
- Keep away from children, food and drinks.

**Antidote:** Atropine, PAM, Toxogonin.

**SAFETY PERIOD:**

- For meat consumption: 14 days.
- For milk consumption: 3 days.

**STORAGE:** Keep in a dry, dark place below 30 °C.

**PACKING:** (100,250, 500, 1000) ML.