

**COMPOSITION:** Each ml contains:

Diclofenac Na 25 m

## **PROPERTIES:**

■ An non-steroidal anti-inflammatory agent of the phenylacetic acid derivative.

Has a powerful anti-inflammatory, analgesic, antipyretic and antispasmolytic actions.

# **INDICATIONS:**

- For the inflammatory and degenerative rheumatic diseases and treatment of painful conditions due to inflammation of non-rheumatic origin.
- Treatment of rheumatoid arthritis, osteo-arthritis, myositis, fibrositis, tendosynovitis and neuritis.
- Soft tissues injuries (Sprain, Traumatic and surgical pain).
- Febrile cases.
- Musculo-skeletal pain.
- Alleviation of swelling in peracute, acute mastitis, metritis, traumatic and surgical wounds.

### **DOSAGE:** Given by intramuscular injection.

- Cattle, Camels, Horses: 4-10 ml per 50 kg B.wt. / day.
- Sheep, Goats, Swine: 0.8-2.4 ml per 10 kg B.wt. / day.
- Dogs, Cats: 0.5-1 ml per 5 kg B.wt. / day.
- Repeat dosage if necessary, but not more than 3-5 days.

### **SAFETY PERIOD:**

- For meat consumption: 28 days.
- For milk consumption: 7 days.

### **REMARKS:**

- Not given in late pregnancy.
- Not given in cases of hepatic, renal disorders or haemorrhages.
- Not given in cases of hypersensitive or allergic animal.
- Transient local action may occur at the injection site.

**STORAGE:** Keep in dry, dark place at 15-25 °C.

**PACKING:** (50, 100) ml.