

DOXYVAP-20%

Water Soluble Powder

COMPOSITION: Each gm contains:

Doxycycline HCl	200 mg
-----------------	--------

PROPERTIES:

- Doxycycline, the broad spectrum antibiotic, is one of the tetracycline group; which is more rapidly absorbed in the gastrointestinal tract, distributed extensively throughout the body, and slowly excreted, than other tetracyclines.
- Doxycycline is a long acting antibiotic, and is a safe treatment even with impaired renal function. It is rapidly absorbed from the intestine and slowly excreted.
- Doxycycline is used in cases of different infections of respiratory, urogenital and gastrointestinal tracts; which are caused by mycoplasma and various gm+ve and gm-ve bacteria (Staph, Strept, Coli, Chlamydia, Clostridia, Rickettsia, Vibrio, Campylobacter, Pasteurella, Leptospira, Spirochaeta, Pseudomonas, Listeria, Shigella, Klesbsiella etc...)

INDICATIONS:

- Given to poultry, calves, lambs, kids, dogs, cats, pigeons and caged-birds.
- For the treatment and prevention of the different diseases: CRD, Coryza, Mycoplasmosis, Air succulitis, Sinusitis, Synovitis, Colibacillosis, Omphalitis and yolk sac infection, Fowl cholera, Necrotic enteritis (Clostridia), and different bacterial enteritis and diarrhea.

DOSAGE: (Treatment duration: 4-5 days)

■ **Treatment:**

Poultry: 100 gm / 200 litres of drinking water.

Pigeons, Caged birds: 0.65 gm / litre of drinking water.

Different animals and pigs: 0.5gm / 10 kg B.wt. daily and (orally with water).

- #### ■ **Prevention:** Half dosage for the same period.

■ **SAFETY PERIOD:**

- For meat consumption : 7 days.
- For eggs consumption: 3 weeks.

- #### ■ **REMARKS:** During treatment, minerals salts should not be added to the medicated water. Not used for table – eggs layers During treatment, eggs not used for human consumption.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.