

**Composition:** Each Gm contains:

Flumequine	200	mg
Colistin sulphate	800,000	I.U.

# **PROPERTIES:**

- Flumequin, a bactericidal antibiotic of the quinolone group, exerting its action in the alimentary and respiratory tracts, beside its rapid absorption and diffusion with quick maximum serum level.
- Colistin (Polymixin group) due to its poor and slow absorption from the intestine, it concentrates its effect locally and thus it shows a remarkable effectiveness against enteric infections. It induces permeability changes within the bacterial cell that result in its death.

# **INDICATIONS:**

## Poultry:

- Cases of CRD, Coryza, Sinusitis, Air Succulitis, Synovitis, Salmonellosis Pasteurellosis, Vibrionic Hepatitis, Colibacillosis, Clostridia, Staph, Enteritis, Diarrhea, Omphalitis and Yolk Sac infection.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

## Calves, Lambs, Kids, Rabbits:

Septicaemia, Enteritis, Diarrhea, Bloody scour, Nephritis, Cystitis etc..

### **DOSAGE:**

(Firstly, dissolve well 100 gm in 1-2 litres of water, then add to the water bulk to be medicated ).

- Treatment:
  - **Poultry**: 100 gm per 200 litres of drinking water for 3-5 days.
  - Calves, Lambs, Kids, Rabbits: 0.25-0.50 gm / 10 kg B.wt. daily and orally for 5 days.
  - **Prevention:** Half dosage for the same period.

**SAFETY PERIOD:** For meat consumption: 7 days.

## **REMARK:**

Trimethoprim – sulphonamide and nitrofurans products should not be associated with flumequine.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.