

LEVOZAN

Oral Liquid

Anthelmintic for sheep, goats, cattle

COMPOSITION: Each ml contains:

Levamisole HCl	30 mg
Oxyclozanide	60 mg

PROPERTIES:

- Oxyclozanide removes practically all flukes (*Fasciola* spp.) present in the bile ducts.
- Levamisole is the racemic mixture of optical isomers of tetramisole .
 - * Levamisole removes mature and developing immature stages of wide range of nematodes, and is highly effective against the following species:
 - **In the abomasum:** *Haemonchus*, *Ostertagia* and *Trichostrongylus* spp.
 - **In the intestines:** *Trichostrongylus*, *Cooperia*, *Nematodirus*, *Oesophagostomum*, *Chabertia* and *Bunostomum*.
 - **In the lungs:** *Dictyocaulus* spp., including inhibited *D. viviparus* larvae.

INDICATIONS:

For the treatment and control of parasitic gastro-enteritis, verminous bronchitis, and liver flukes infestation.

- **Gastro-intestinal worms:** *Nematodirus*, *Haemonchus*, *Ostertagia*, *Cooperia*, *Trichostrongylus*, *Bunostomum*, *Chabertia*, *Strongyloides*, *Oesophagostomum*.
- **Lung worms:** *Dictyocaulus* spp.
- **Tape worms:** *Moniezia* spp.
- **Liver flukes:** *Fasciola* spp.

DOSAGE: Shake well before use.

Sheep, Goats, Cattle: 2.5 ml per 10 kg B. wt.

(7.5 mg levamisole HCl / kg B. wt. , 15 mg oxyclozanide / kg B. wt.)

SAFETY PERIOD:

- For meat consumption: 18 days.
- For milk consumption: 3 days.

REMARKS:

- Can be given to young, pregnant and lactating animals.
- Animals can be treated with organophosphorous insecticides after 14 days of the anthelmintic drenching.

STORAGE:

- Keep in a dry, dark place at below 30 °C.
- Keep out of reach of children.

PACKING: (100, 250, 500, 1000) ML.