

# ***OXYDONE-FORTE 20% LA***

## **INJECTABLE SOLUTION**

**For livestock**

**COMPOSITION:** Each ml contains:

Oxytetracycline (As dehydrate)

200 mg

## **PROPERTIES:**

- A broad spectrum antibiotic against susceptible Gm+ve bacteria (Staph, Strept, Coryne, Clostridia, Erysipelothrix) and Gm-ve bacteria (E.coli, Pasteurella, Actinobacillus, Moraxella), Mycoplasma, Protozoa, Haemophilus, Spirochaetes, Chlamydia, Leptospira, Rickettsia, Listeria and some large viruses.
- Provides therapeutic effectiveness of up to (4) days duration following a single treatment.
- Well tolerated and distributed rapidly and extensively throughout the body tissues, reaching its peak level in short time and remains active for long period.

## **INDICATIONS:**

- For Sheep, Goats, Cattle, Buffaloes, Camels, Pigs.
- Treatment of different cases:
  - Shipping fever, Wooden Tongue (Actinobacillus), Pink eye (Bovine and ovine kerato conjunctivitis "Moraxella"), Swine Dysentery (Spirochaetosis), Atrophic rhinitis in pigs, Dysagalactia, Foot rot, Calf diphtheria.
  - Anaplasmosis (Theileriosis) and Heart water in cattle (Ehrlichia "Rickettsia"), Red water of calves (Leptospira).
  - Enzootic abortion (Chlamydiosis) in sheep and goats.
  - Clostridial infections: Enterotoxaemia (Lamb dysentery, Black leg, Malignant oedema).
  - Scours, enteritis, diarrhea.
  - Septicaemia, Pyaemia, Peritonitis, Navel / Joint-ill, wounds and skin infections, abscesses.
  - Supportive therapy in mastitis.
  - Respiratory tract infections: Bronchitis, pneumonia, rhinitis, etc.....
  - Urinary tract infections: Nephritis, Pyelonephritis, Cystitis.
  - Genital tract infections: Metritis, Puerperal Sepsis, etc...
  - Control of post-operative and post-parturient infections.
  - Bacterial complications of viral diseases.

**DOSAGE:** By deep intramuscular injection.

- Cattle, Buffaloes, Camels, Sheep, Goats, Pigs: 1 ml per 10 kg B.wt. in one single dose.
- In severe cases, repeat the injection dose after 3-5 days.
- Inject no more than (20 ml) in cattle, buffaloes, camels and no more than (5 ml) in sheep, goats, pigs to be given at the same site.

## **SAFETY PERIOD:**

- For meat consumption: 21 days.
- For milk consumption: 7 days.

## **REMARKS:**

- Not recommended for horses, dogs, cats.
- Divide the dosage into separate injection sites to minimize local tissue irritation.
- Transient localized swelling may occur at the site of injection for few days.
- Occasional allergic reaction may occur, in such cases, discontinue use and give antihistaminic (epinephrine etc....) immediately, beside symptomatic treatment, if necessary.
- Not given with penicillins, calcium and iron preparations as interference action may occur.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** ( 50, 100) ml.