

# ***RENAMIDE***

**injectable solution  
diuretic**

**COMPOSITION** : Each ml contains :

Furosemide

50 mg

**PROPERTIES** :

- potent diuretic with rapid onset action
- increased na and water excretion without loss of potassium.
- effective even in cases of glomerular – infiltration impairment .

**INDICATIONS** :

- for cattle , equines, dogs, cats, pigs.
- pathological oedema of the udder and perineum of the cow and mare at time of parturition .
- preputial or scrotal oedema of male animals .
- supportive treatment in the elimination of fluids in cases of ascites, hydrothorax , and bronchopneumonia , hydropericardium , from joints and tendon sheaths of bursae .
- supportive treatment in laminitis, paralytic myoglobinuria of the horse .
- protein deficiency oedema due to renal proteinuria , liver lesions caused by protein metabolism disorders, parasitic infestations or mal- nutrition seen as generalised skin oedema and ascites .
- skin oedema due to blood vessels lesions of toxicity, trauma, allergy , inflammations , wounds, surgery.

**DOSAGE** : ( By Intramuscular Or Slow Intravenous Injection )

- **Equines** : 5-10 ml once or twice daily with intervals of 6-8 hours.
- **Cattle** : 10 ml one time, or 5 ml twice with interval of 6-8 hours .
- **Dogs ,Cats**: 0.25-0.5 ml /5 kg b.wt. once or twice daily with interval of 6-8 hours.
- **Pigs** : 1 ml/10 kg b.wt once or to be determined empirically.

**SAFETY PERIOD** :

for meat and milk consumption : 2 days .

**REMARKS** :

- ◀ not used in cases of renal failure with anuria, oliguria, acute glomerular nephritis , electrolytes deficiency disease or over dosage with digitalis .
- ◀ the therapeutic effect may be impaired by increased water intake .
- ◀ diuresis may be enhanced with oral nacl intake.
- ◀ in hypocalcaemic animals , tetany may occur.

**STORAGE** :

Keep in dry, dark place below 30 °C.

**PACKING** : (50,100) ML .