<u>TETRAMISOLE – 30</u>

Water Soluble & Feed additive Powder The safe & broad spectrum vermicidal, larvicidal anthelmintic for livestock

COMPOSITION: Each gm contains:

Tetramisole HCl 300 mg

PROPERTIES:

- Belongs to the imidothiazole group of anthelmintic which is very effective against nematodes
- It is the racimic form of levamisole HCl.

INDICATIONS:

For the treatment and control of parasitic gastro – eneritis and verminous bronchitis caused by round worms (Nematodes):

- Ruminants, Rabbits:
 - Gastro intestinal worms: Ascaris, Nematodirus, Haemonchus, Ostertagia, Cooperia, Thrichuris, Strongyloides.
 - Lung worms: Dictyocaulus.
- **Poultry:** Ascaridia, Heterakis, Capillaria, Amidostomum.

DOSAGE:

- Given with water or feed.
- Give single dose each time.
- Sheep, goats, cattle, camels: 0.5 gm per 10 kg B. wt. (15 mg active sub. / kg body wt.).

Repeat for prevention every 4-6 months.

- Poultry, Pigeons: 0.06 gm per kg B. wt. (20 mg active sub. / kg body wt.) Repeat every 3 months.
- **Rabbits:**0.06 gm per kg B. wt. (20 mg active sub. / kg B. wt.) Repeat twice per year.

SAFETY PERIOD:

- For meat and eggs consumption: 7 days.
- For milk consumption: 1 day.

REMARK: Organophosphates insecticides are toxic if applied within (14) days of treatment. **STORAGE:** Keep in a dry, dark place at below 30 °C. **PACKING:** (100, 250, 500, 1000) GM.