

<u>COMPOSITION</u> : each ml contains :

tilmicosin 300 mg

PROPERTIES :

- a long acting macrolide antibiotic which is effective against gm +ve and some gm ve bacteria ,and mycoplasmas as well .
- maintains a high therapeutic effect for about three days after dosing .

INDICATIONS :

for the treatment of bovine respiratory diseases caused by pasteurella multocida , manheimia (pasteurella) haemolytica , haemophilus somnus, histophilus somni and mycoplasma spp.

DOSAGE :

- by a single subcutaneous injection in the neck or behind the shoulder . 1 ml per 30 kg b.wt.
 - (10 mg activity / kg b.wt)
- do not overdose.
- not used in calves less than 70 kg b.wt.

SAFETY PERIOD :

for meat consumption : 28 days .

REMARKS :

- not recommended for goats, horses and pigs.
- the dose should be subdivided .
- do not inject more than 15 ml per injection site.
- transient localized swelling may occur at the injection site.
- do not use in milking animals .

PRECAUTIONS:

- before dosing, wear protective gloves and safety goggles.
- avoid contact with eyes.
- wash hands with soap before eating , drinking or smoking.

WARNING:

strictly, avoid accidental self injecting, and in such case, seek immediate medical attention, and apply ice or cold pack on the injection site.

STORAGE :

- keep out of reach of children.
- Keep in dry, dark place below 30 °C.

PACKING: (50, 100) ml.