

# **TRIPLE SULPHA / TRIPLE SULPHA – S**

## **Water Soluble Powder**

**COMPOSITION:** Each gm contains

	<b>TRIPLE SULPHA</b>	<b>TRIPLE SUIPHA - S</b>
SULPHADIMIDINE Na	250 MG	125 MG
SULPHADIAZINE Na	250 MG	125 MG
SULPHATHIAZOLE Na	250 MG	125 MG

### **PROPERTIES:**

- 3-Combined valuable sulpha components, with a high antibacterial activity against Gm +ve and Gm –ve bacteria: Coli, Staph, Strept, Salmonella, Pasteurella, Proteus, Chamydia etc..
- Also effective against Eimeria spp. (Coccidia).
- Sulphonamides diffuse very widely into tissues and thus exert effectiveness against the bacterial infections.
- Sulphonamides are well tolerated and have low toxicity, hence it is safe to be used.

### **INDICATIONS:**

- **Poultry:** Coryza, Caecal coccidiosis.
- **Ruminants:** Pneumonia, Enteritis, Scour, Foot rot, Haemorrhagic septicaemia (Pasteurella)
- **Equines:** Pneumonia, Strangles, Enteritis, Diarrhea.

### **DOSAGE:**

- **Treatment:** Duration for 3-5 days.

	<b>TRIPLE SULPHA</b>	<b>TRIPLE SUIPHA - S</b>
<b>Poultry</b>	250 gm per 200 litres of drinking water	500 gm per 200 litres of drinking water
<b>Ruminants, Equines</b>	10-15 gm per 50 kg B. wt. daily and orally with water	20-30 gm per 50 kg B. wt. daily and orally with water

- **PREVENTION:** Half dosage for 3 days.

### **SAFETY PERIOD:**

- For meat consumption: 7 days.
- For milk consumption: 4 days.

### **REMARKS:**

- Not recommended for young stocks and layers.
- Ensure adequate water intake during treatment to reduce the risk of crystalluria.

**STORAGE:** Keep in a dry, dark place below 30 °C.

**PACKING:** (100, 250, 500, 1000) GM.

