

COMPOSITION: Each gm contains: Tylosin tartrate (activity) Doxycycline HCl

200 mg 200 mg

## **PROPERTIES:**

- A combination of two broad specturm antibiotics which are effective against different Gm +ve and Gm -ve bacteria and some large viruses.
- Tylosin is very active against respiratory infections.
- Doxycycline, the rapid absorption and slow excretion of doxycycline exerts additional activity.

# **INDICATIONS:**

- Prevention and treatment of different infections of the respiratory, alimentary and urogenital tracts such as: Mycoplasmosis, CRD, Coryza, Colibacillosis, Omphalitis and Yolk sac infection, Enteritis and diarrhea, Pasteurellosis, Necrotic enteritis, Air succulitis, Sinusitis, Synovitis, Staph, Strept, Coryne, Hexamitiasis, Spirochaeta.
- Stress conditions due to vaccination, rehousing, debeaking, transporting, changes of feed and weather.

#### DOSAGE: (Treatment duration: 4-5 days).

- Poultry:
  - **Treatment:** 100 gm per 200 litres of drinking water.
  - **Prevention:** Half dosage for the same period.
- Different animals: 0.5 gm per 10 kg B.wt. daily and orally with water.

## **SAFETY PERIOD:**

For meat consumption: 7 days.

## **REMARK:**

- Not recommended for table-eggs layers.
- During application, do not medicate water with minerals salts.

**STORAGE:** Keep in a dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.