

# **VAPCOTRIL – 10**

## **Oral Solution**

**COMPOSITON:** Each ml contains:

Norfloxacin                    100 mg

### **PROPERTIES:**

- Characterized by quick absorption and rapid high serum level.
- Has a broad spectrum efficacy against a wide range of bacteria and mycoplasmas:
  - Gm+ve bacteria: Staph, Strept, Coryne, Clostridia and Listeria.
  - Gm-ve bacteria: E. coli, Slamonella, Klebsiella, Proteus, Yersina, Vibrio, Haemophilus, Pasteurella, Actinobacillus, Pseudomonas, Bordetella, Brucella, Moraxella and Campylobacter.
  - Mycoplasmas: M. gallisepticum, M. synoviae, M. meleagridis.....etc.

### **INDICATIONS :**

For prevention and treatment of different infections:

- **Poultry:** CRD, Colibacillosis, Coryza, Fowl cholera, Fowl typhoid, other Salmonellosis (S. typhimurium, S. enteritidis etc..), Staph and Strept infections, Mycoplasmosis, Hexamitiasis Vibrionic hepatitis etc...
- **Calves, Lambs:** Enteritis, diarrhea and respiratory infections of newly-born and growing off-springs due to Colibacillosis, Pasteurellosis, Salmonellosis Mycoplasmosis, and Clostridia, Vibrionic dysentery and Synovitis as well.
- **Rabbits:** Pasteurellosis, Enterotoxaemia, enteritis, dysentery, snuffle, respiratory infections, Listeriosis, endometritis, Tyzzer's disease and Yersinia infection (Rodentiose).

**DOSAGE:** ( Treatment duration for 3-5 days ).

- **Treatment:**
  - **Poultry:** 100 ml per 200 litres of drinking water.
  - **Calves, Lambs, Rabbits:**  
0.3-0.5 ml / 10 kg B.wt. daily and orally with water.  
(3-5 mg activity per kg B.wt. daily)

**Prevention:** Half dosage for the same period.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** (100, 250, 500, 1000) ML.