

# ***DEWORMER – 5%***

**Oral Suspension**

**Anthelmintic for Sheep, Goats, Cattle, Camels, Equines.**

**COMPOSITION:** Each ml contains:

**Fenbendazole**

**50 mg**

## **PROPERTIES:**

- An effective, safe and broad spectrum anthelmintic of the benzimidazole group.
- It has vermifugal, larvicidal and ovicidal activity against nematodes (Round Worms) and cestodes (Tape worms).

## **INDICATIONS:**

For treatment and control of parasitic gastro – enteritis and verminous bronchitis.

- **Sheep, Goats, Cattle, Camels:**
  - **Gastrointestinal Round Worms:** Ostertagia, Trichostrongylus, Haemonchus, Strongyloides, Nematodirus, Cooperia, Bunostomum, Oesophagostomum, Marshallagia, Chabertia...
  - **Lung Worms:** Dictyocaulus, Muellerius.
  - **Tape Worms:** Moniezia.
- **Equines:** Small and large Strongyles, Oxyuris, Ascarids, Trichostrongylus, Strongyloides.

**DOSAGE:** (single dose).

- **Shake well before use.**
- **Sheep, Goats:** 1 ml / 10 kg B. wt. (5 mg / kg B. wt.)
- **Cattle, Camels:** 1.5 – 2 ml / 10 kg B. wt. (7.5 mg / kg B. wt.)  
For tape worms in cattle: 2 ml / 10 kg B. wt.  
(10 mg activity / kg B. wt.)
- **Equines:** 1-2 ml / 10 kg B. wt. (5mg activity / kg B. wt.)  
For ascarids in equines: 2 ml / 10 kg B. wt.  
(10 mg / kg B. wt.)

## **SAFETY PERIOD:**

- For meat consumption: 14 days.
- For Milk consumption: 3 days.

## **REMARKS:**

- Repeat treatment after 4-6 weeks, when necessary.
- Regular deworming at intervals of 6-8 weeks may be required for equines.

**STORAGE:** Keep in a dry, dark place at below 30 °C .

**PACKING:** (100, 250, 500, 1000) ML.