

# DIAZINON-60%

**PUBLIC HEALTH INSECTICIDE  
EMULSIFIABLE CONCENTRATE**

**Class :  
Organophosphate**

Public health insecticide with contact, stomach and respiratory action to control insects that affect public health such as: Cockroaches, Housefly, Mosquitoes, Bugs, Fleas and Ants,...etc .

**COMPOSITION :** Each liter contains:

**DIAZINON**

**60% W/V (A.I.)**

**PROPERTIES:**

**DIAZINON-60%:** An Organophosphate insecticide acts by contact, stomach and respiratory action.

**DIAZINON-60%:** Used to control public health insects such as: Cockroaches, Bedbugs, Ants, Lice, Fly larvae, House flies, Fleas and Ticks.

**DIAZINON-60%:** Has long residual effect.

**USAGE AND DOSAGE:**

PEST	APPLICATION RATES	NOTICE
<b>* SURFACE SPRAY</b>		
Cockroaches	100-200 ml / 20 lt. water/ 100 m <sup>2</sup>	Spray infected areas such as baseboards, areas around water pipes, underneath sinks, cracks, crevices, surfaces behind and beneath cabinets, refrigerators and other places where insects hide. Repeat spraying every 1-2 months to avoid re-infection.
flies and Mosquitoes		Spray walls and ceilings indoors and other surfaces on which flies and mosquitoes resting places out-side the building such as verandas, eaves, outhouses..etc.
Bedbug And Lice		Spray walls and bed frames, and under sides of mattresses as well as stuffed furniture. Pay particular attention to cracks, crevices, back of picture, windows and door frames.
Silverfish		----
Ants	150 ml /20 lt. water/ 100 m <sup>2</sup> (you can add 100 gm sugar to 4-5 lt. of spray mixture).	Wet ant hills and runways, apply at house entrances Door and window frames, cracks in walls.
<b>* SPACE TREATMENT (THERMAL FOGGING OR ULV):</b>		
Mosquitoes	330-660 ml / 2-10 lt. diesel /ha.	Space treatments against adult Mosquitoes should be carried out Regularly in streets, market places, in parks, playgrounds, beaches, rubbish tips.
Flies		Space treatments against adult Flies should be carried out regularly in streets, market places, in parks, playgrounds, beaches, rubbish tips.
<b>* USING PRESSURE SPRAYER OR WATERING CAN:</b>		
Fly larvae	30 ml /10 lt. water.	Restrict treatment against Fly larvae to cases where breeding ground are localized and clearly defined (manure heaps, open latrines). Breeding sites should be soaked with spray liquid. Apply spray mixture as a coarse spray or with a watering can. Application should be repeated at weekly intervals.

Hectare = 10000 m<sup>2</sup>

**RE-ENTRY PERIOD:** 48 hours after the last treatment.

**RESIDUAL ACTIVITY:** About 4-6 weeks.

**NOTICE :** Avoid spraying **DIAZINON-60% (PH)** when weather temperature below -10 °C and above + 35 °C and away from humidity.

- For more details about first aid & precautions please refer to first aid & precautions index .