

FLORAM – 30%

injectable solution

COMPOSITION : each ml contains :

florfenicol 300 mg

PROPERTIES :

- florfenicol is highly active and broad spectrum antibiotic against many gm +ve and gm-ve bacteria .
- it acts by binding the ribosomal subunit and inhibiting the bacterial protein synthesis.

INDICATIONS :

for the treatment of the bacterial infections in cattle and pigs.

- cattle :
 - the respiratory diseases caused by the
 - bacteria : pasteurella multocida , mannheimia haemolytica , histophilus somni.
 - foot rot caused by the bacteria : fusobacterium necroforum , bacteroides melaninogenicus .
- pigs : the respiratory diseases caused by the bacteria : actinobacillus pleuropneumoniae, pasteurella multocida, salmonella cholerae suis , streptococcus suis .

DOSAGE : (by intramuscular or subcutaneous injection)

- cattle : (in the neck muscle)
 - 10 ml / 150 kg b.wt.
 - (20 mg activity / kg b.wt
 - repeat the dose after 2 days in another site of the neck muscle .

OR

 - by a single subcutaneous injection in the neck region :
 - 20 ml / 150 kg b.wt.
 - (40 mg activity / kg b.wt.)
- pigs : by intramuscular injection .
 - 1 ml / 20 kg b.wt.
 - (15 mg activity / kg b.wt.)
 - repeat the dose after 2 days .

SAFETY PERIOD :

for meat consumption :
cattle : 28 days (for i.m. injection)
 38 days (for s.c. injection)
pigs : 18 days

REMARKS :

- the dose should be subdivided .
- do not inject , in each site , more than 10 ml for cattle , and not more than 3 ml for pigs.
- not given to milking cattle .
- not used for animals of breeding age.

STORAGE : Keep in dry, dark place below 30 °C.

PACKING : (50,100) ml .