

OXYDONE-3%

INJECTABLE SOLUTION

For livestock

COMPOSITION: Each ml contains:

Oxytetracycline HCl

30 mg

PROPERTIES:

- A broad spectrum antibiotic against susceptible Gm+ve bacteria (Staph, Strept, Coryne, Clostridia, Erysipelothrix) and Gm-ve bacteria (E.coli, Pasteurella, Actinobacillus, Moraxella), Mycoplasma, Protozoa, Haemophilus, Spirochaetes, Chlamydia, Leptospira, Rickettsia, Listeria and some large viruses.
- Distributed rapidly and extensively throughout the body tissues.

INDICATIONS:

- For Sheep, Goats, Cattle, Buffaloes, Camels, Pigs, Dogs, Cats, Rabbits, Poultry.
- Treatment of different cases:
 - Shipping fever, Wooden Tongue (Actinobacillus), Pink eye (Bovine and ovine kerato conjunctivitis “Moraxella”), Swine Dysentery (Spirochaetosis), Atrophic Rhinitis in pigs, Dysagalactia, Foot rot, Calf diphtheria.
 - Anaplasmosis (Theileriosis) and Heart water in cattle (Ehrlichia “Rickettsia”), Red water of calves (Leptospira).
 - Enzootic abortion (Chlamydiosis) in sheep and goats.
 - Clostridial infections: Enterotoxaemia (Lamb dysentery, Black leg, Malignant oedema).
 - Scours, enteritis, diarrhea.
 - Septicaemia, Pyaemia, Peritonitis, Navel / Joint-Ill, wounds and skin infections, abscesses.
 - Supportive therapy in mastitis.
 - Respiratory tract infections: Bronchitis, pneumonia, rhinitis, etc.....
 - Urinary tract infections: Nephritis, Pyelonephritis, Cystitis.
 - Genital tract infections: Metritis, Puerperal Sepsis, etc...
 - Control of post-operative and post-parturient infections.
 - Bacterial complications of viral diseases.
 - Poultry: Colibacillosis, Sinusitis, Coryza, CRD, Blue Comb, Mycoplasmosis.
 - Rabbits: Diarrhea, Pneumonia, Enteritis, etc....

DOSAGE: (By deep intramuscular, subcutaneous injection).

Treatment duration: 3-5 days.

- Cattle, Buffaloes, Camels, Sheep, Goats, Pigs: 0.5-1.3 ml per 10 kg B.wt. daily.
- Dogs, Cats: 0.8-1.6 ml per 5 kg B.wt. daily.
- Rabbits, Poultry: 0.8-3.2 ml per animal daily.
- Divide the dosage into separate injection sites to minimize the local tissue irritation.

SAFETY PERIOD:

- For meat consumption: 14 days.
- For milk consumption: 5 days.

REMARKS:

- Transient localized swelling may occur at the site of injection for few days.
- Not given with penicillins, calcium and iron preparations as interference action may occur.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (50,100) ml .