

OXYDONE-FORTE 30% LA

**injectable solution
for livestock**

COMPOSITION: Each ml contains:

Oxytetracycline (As dihydrate) 300 mg

PROPERTIES:

- A broad spectrum antibiotic against susceptible Gm+ve bacteria (Staph, Strept, Coryne, Clostridia, Erysipelothrix) and Gm-ve bacteria (E.coli, Pasteurella, Actinobacillus, Moraxella), Mycoplasma, Protozoa, Haemophilus, Spirochaetes, Chlamydia, Leptospira, Rickettsia, Listeria and some large viruses.
- Provides therapeutic effectiveness of up to (4) days duration following a single treatment.
- Well tolerated and distributed rapidly and extensively throughout the body tissues, reaching its peak level in short time and remains active for long period.

INDICATIONS:

- For Sheep, Goats, Cattle, Buffaloes, Camels, Pigs.
- Treatment of different cases:
 - Shipping fever, Wooden Tongue (Actinobacillus), Pink eye (Bovine and ovine kerato conjunctivitis “Moraxella”), Swine Dysentery (Spirochaetosis), Atrophic rhinitis in pigs, Dysagalactia, Foot rot, Calf diphtheria.
 - Anaplasmosis (Theileriosis) and Heart water in cattle (Ehrlichia “Rickettsia”), Red water of calves (Leptospira).
 - Enzootic abortion (Chlamydiosis) in sheep and goats.
 - Clostridial infections: Enterotoxaemia (Lamb dysentery, Black leg, Malignant oedema).
 - Scours, enteritis, diarrhea.
 - Septicaemia, Pyaemia, Peritonitis, Navel / Joint-Ill, wounds and skin infections, abscesses.
 - Supportive therapy in mastitis.
 - Respiratory tract infections: Bronchitis, pneumonia, rhinitis, etc.....
 - Urinary tract infections: Nephritis, Pyelonephritis, Cystitis.
 - Genital tract infections: Metritis, Puerperal Sepsis, etc...
 - Control of post-operative and post-parturient infections.
 - Bacterial complications of viral diseases.

DOSAGE: By deep intramuscular injection.

- Cattle, Buffaloes, Camels: 0.7 ml per 10 kg B.wt. in one single dose.
- Sheep, Goats, Pigs: 0.3-0.7 ml per 10 kg B.Wt.
- In severe cases, repeat the injection dose after 3-5 days.
- Inject no more than (20 ml) in cattle, buffaloes, camels and no more than (5 ml) in sheep, goats, pigs to be given at the same site.

SAFETY PERIOD:

- For meat consumption: 28 days.

- For milk consumption: 7 days.

REMARKS:

- Not recommended for horses, dogs, cats.
- Divide the dosage into separate injection sites to minimize local tissue irritation.
- Transient localized swelling may occur at the site of injection for few days.
- Occasional allergic reaction may occur, in such cases, discontinue use and give antihistaminic (epinephrine etc....) immediately, beside symptomatic treatment, if necessary.
- Not given with penicillins, calcium and iron preparations as interference action may occur.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (50, 100) ml.