

TYLOSIN-20%

Oral Solution
For Poultry, Turkey, Pigeons

COMPOSITION: Each ml contains:

Tylosin Tartrate (activity) 200 mg

PROPERTIES:

- Tylosin as a strong macrolide antibiotic covers a wide spectrum of Gm +ve bacteria such as: Staph, Strept, Diplococcus, Coryne, Clostridia and other bacterial infections secondary to viral diseases.
- Very effective against mycoplasma infections and affects also some rickettsias (Group psittacosis-chlamydiosis) and Haemophilus gallisepticum.
- Some Gm –ve bacteria (Pasteurella, Campylobacter) are also affected by tylosin.

INDICATIONS:

- For prevention and treatment of: Mycoplasmosis, CRD, Air succulitis, Infectious Synovitis, Infectious Sinusitis, Chlamydiosis, Spirochaetosis.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

DOSAGE:

- Respiratory or general infections in poultry:
500 ml per 200 litres of water for 3-5 days according to severity of the disease.
- Spirochaetosis: (2.5% solution), used as preventive and for control.
- Pigeons: 0.075-0.125 ml of the product / kg B.wt. / 4 times daily.
(15-25 mg activity / kg B.wt.)

PREVENTION PROGRAM:

Use 500 ml per 200 litres of water as follows:

- **Broilers:** 1st week of age for 3 days.
4th week of age for 1 day.
- **Pullets:** 1st week of age for 3 days.
4th week of age for 1 day.
9th week of age for 2 days.
20th week of age for 2 days.
- **Parent stocks:** 1st week of age for 5 days.
4th, 9th, 16th, 20th and 24th week of age for 2 days.
- **Turkeys:** 1st week of age for 5 days.
3rd week of age for 1 day.

REMARK:

- Not recommended for table-eggs layers.

SAFETY PERIOD: For meat consumption: 7 days.

STORAGE: Keep in a dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) ML.