

***VAPCOTRIM-PLUS / VAPCOTRIM-PLUS Potentiated***  
**Water Dispersible Powder**  
**For Poultry, Turkeys & other Animals**

**COMPOSITION:** Each gm contains:

	<b>VAPCOTRIM-PLUS</b>	<b>VAPCOTRIM-PLUS/POTENTIATED</b>
	-----	-----
Erythromycin Thiocyanate	80 mg	180 mg
Trimethoprim	30 mg	30 mg
Sulphadiazine-Na	150 mg	150 mg

**PROPERTIES:**

- Erythromycin inhibits the protein synthesis of bacteria, while trimethoprim magnifies the effect of sulphonamide by blocking transformation of folic acid into bio-active folic acid.
- Effective against Gm+ve and Gm-ve bacteria: Coli, Staph, Strept, Salmonella, Pasteurella, Vibrio, Mycoplasma, Clostridia, Listeria, Coryne etc...

**INDICATIONS:**

- **Poultry:**
  - For prevention and treatment of: CRD, Mycoplasmosis, Coryza, Infectious sinusitis, Air succulitis, Infectious synovitis, Vibronic hepatitis, Salmonellosis, Pasteurellosis, Colibacillosis, enteritis, diarrhea.
  - Stress conditions due to vaccination, debeaking, rehousing, transporting, changes of feed and weather.
- **Other animals:** Cases of pneumonia, bronchitis, enteritis and diarrhea etc..

**DOSAGE:**

- **Treatment:** (Duration for 5 days).
  - **Poultry:** 100 gm per 200 litres of drinking water.
  - **Other animals:** 2 gm per 15 kg B.wt. daily and orally with water.
- **Prevention:** Half dosage for the same period.

**REMARKS:**

- Flumequine should not be associated with trimethoprim-sulphonamide treatment.
- Not recommended for laying flocks.
- Contra-indicated in cases of liver and Kidney troubles, Gumboro disease.

**SAFETY PERIOD:** For meat and milk consumption: 7 days.

**STORAGE:** Keep in a dry, dark place below 30 °C.

**PACKING:** (100, 250, 500) KG.