

# ***VIROSEPT***

## **Solution**

### **The Strong Disinfectant and Detergent**

#### **COMPOSITION:**

Cetrimide	6 %
Glutaraldehyde	6 %
Formaldehyde	8.4 %

#### **PROPERTIES:**

- A powerful combination of 3 effective and synergistic disinfectants.
- It is characterized by strong detergent and disinfecting activity against the bacteria and spores, viruses, fungi, moulds and algae.
- Has a broad spectrum action against all pathogenic micro organisms such as viruses (Avian Influenza, Swine influenza, ND, IB, 1LT, IBD etc...).
- Has a long lasting action which prevents recontamination.

#### **APPLICATIONS:** (Before spraying, dilute with water 1:200 – 1:400 (0.25 - 0.50%).)

- **General disinfection:**  
Farm buildings, poultry and animals houses, battery systems.  
Leave for 12 hrs. contact.
- **Farm equipment, trucks, vehicles:** Leave for 30 minutes contact.
- **Hatcheries (Incubators, hatchers, equipment, floors):** Leave for 60 minutes contact.
- **Footbaths.**
- **Fog device:** Ambient disinfection (1.5 litres virosept + 3 litres water) per 1000 cubic meters volume.
- For area spraying, use 1 litre of the diluted product per 4 sq. meters.

#### **FIRST AID:**

- **Eyes:** Flush with water for at least 15 minutes.
- **Skin:** Wash with soap and water.
- **Inhalation:** Remove to fresh air ,apply artificial respiration or give oxygen .
- **Ingestion:** Do not induce vomiting and do not give anything to drink.
- **Get medical attention when necessary.**
- **Antidote:** Not specific, treat according to symptoms.

#### **PRECAUTIONS:**

- Keep away from children, food and drinks.
- Avoid contact with eyes, mouth, skin and inhalation.
- Do not pollute ponds, water sources and canals.
- Before application, wear protective clothes, gloves, boots, goggles.
- Do not eat, drink or smoke during spraying.
- Wash containers with water before discarding.

#### **STORAGE:** Keep in a dry, dark place at below 30 °C.

#### **PACKING:** (100, 250, 500, 1000) ML. 5 litres gallon.