

ADIFON-25%

**FUNGICIDE
WETTABLE POWDER**

VAPCO

Class:
Triazole

Systemic fungicide used to control Powdery mildew, Rusts, Black rot, Leaf blotch and Leaf spots on crops mentioned in the table of uses.

COMPOSITION: Each kg contains:

TRIADIMEFON

25% W/W (A.I)

PROPERTIES:

ADIFON-25% : Systemic fungicide with protective and curative action absorbed by roots and leaves and translocated rapidly in young growing tissues.

ADIFON-25% : Control many fungal diseases on crops mentioned in the table of uses.

ADIFON-25% : Slightly toxic to mammals.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE	SAFETY PERIOD (DAYS)
Grapes	Black Rot, Powdery Mildew	150 - 200 Gm/ha	14
Tomatoes, Pepper and Artichokes,	Powdery Mildew	500 Gm/ha	14
Eggplant, Cucumber, Onion, Turnip	Black Rot, Powdery Mildew	400 Gm/ha	14
Okra	Black Rot, Powdery Mildew	238 Gm/ha	7
Sugar Beet	Powdery Mildew, Beet Rust	500 Gm/ha	14
Cereals	Powdery Mildew, Rust, Leaf Blotch, Leaf Spot.	500 - 1000 Gm/ha (Repeat The Treatment When Reinfestation Is Observed Again)	28
Peas	Powdery Mildew	400 Gm/ha	14
Beans	Rust	500 Gm/ha	14
Tobacco	Powdery Mildew	10 - 20 Gm/100 lt. Water (Repeat Spray Every 7 - 14 Days)	14
Coffee	Rust	0.5 - 1 Kg/ha (Protective spray or when first symptoms of disease appear). 1 - 2 Kg/ha (in case of severe rust infection repeat every 3 - 4 weeks).	3 - 4 Weeks

Hectar= 10000 m²

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY :

Compatible with most pesticides, spray mixture must be applied immediately after being prepared.

• For more details about first aid & precautions please refer to first aid & precautions index.