

AGRINATE-25%

INSECTICIDE
WETTABLE POWDER

VAPCO

Class:
Carbamate

Systemic insecticide used to control a wide range of agricultural crops.

COMPOSITION: Each kg contains:

METHOMYL

25% W/W (A.I)

PROPERTIES:

AGRINATE-25%: Packed in water soluble sachets.

AGRINATE-25%: Has a fast knockdown on a wide range of insects.

AGRINATE-25%: Has a broad spectrum activity against a wide range of insects at all developing stages (eggs, larvae, adults).

AGRINATE-25%: Is extremely active against the most difficult insect pests such as Borer, Fruit worm, Leaf miner which have become resistant to Organophosphorus and Pyrethroid compounds.

AGRINATE-25%: Quickly decomposes in nature, which decreases the safety period for Vegetables and Fruit trees.

AGRINATE-25%: Non persistent in the environment.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE	SAFETY PERIOD (DAYS)
Tomatoes	Fruitworm, Leafworm	1 - 2 bags/drum	7
Cabbage. Cauliflower	Green worms, Leafworm, Aphids, Thrips.	1 - 2 bags/drum	7
Cucumber, Squash, Watermelon, Beans, Jew's mallow, Pepper, Eggplant, Potatoes.	Leafworm, Aphids, Thrips.	2 bags/drum	7
Corn	Cutworm, European Corn Borer, Aphids, Beetles.	2 bags/drum	7
Apples, Pear, Peach, Apricot, Cherries, Almonds and Citrus	Green and Black aphids, Leafminer, Stem Borer, Psylla, Leafroller, Thrips, Leafhopper.	2 bags/drum	14
Grapes	Grape berry moth, Leafhopper.	2 bags/drum	7
Ornamentals (Open Field, Green Houses)	Cotton Leafworm, Cutworm, Bugs, Aphids, Thrips, Leafhopper.	2 bags/drum	--

Bag = 200 gm of **AGRINATE-25%**

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

72 hours after the last application.

REMARKS:

- Use sprayers with low volume pressure, and use for large farms and it is not recommended for use in gardens.
- Fill 1/4 or 1/2 of the spray tank with water, then add the recommended dose of **AGRINATE-25%**, stir, fill the tank and start spraying.
- Use the recommended quantities at the beginning of infestation, repeat spraying 5 - 7 days intervals.
- In case there is more than one dose recommended, use the lower dosage for young crops or when the injury is slight.

COMPATIBILITY:

Compatible with most common insecticides and fungicides. It's recommended to make mixing trial before mixing.

• For more details about first aid & precautions please refer to first aid & precautions index.