

AGRINATE-90%

INSECTICIDE
SOLUBLE POWDER

VAPCO

Class:
Carbamate

Systemic insecticide used to control a wide range of agricultural crops.

COMPOSITION: Each kg contains:

METHOMYL

90% W/W (A.I)

PROPERTIES:

AGRINATE-90%: Packed in water soluble sachets.

AGRINATE-90%: Has a fast knockdown on a wide range of insects.

AGRINATE-90%: Has a broad-spectrum activity against a wide range of insects at all developing stages (eggs, larvae, adults).

AGRINATE-90%: Is extremely active against the most difficult insect pests such as Borer, Fruitworm, Leafminer which have become resistant to Organophosphate and Pyrethroid compounds.

AGRINATE-90%: Quickly decomposes in nature, which decreases the safety period for vegetables and fruit trees.

AGRINATE-90%: Non persistent in the environment.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE	SAFETY PERIOD (DAYS)
Tomatoes	Fruitworm, Leafworm	1-2 bag / 2 drums	7
Cabbage, Cauliflower	Green Worms, Leafworm, Aphids, Thrips.	1-2 bag / 2 drums	7
Cucumber, Squash, Watermelon, Beans, Jew's mallow, Pepper, Eggplant, Potatoes.	Leafworm, Aphids, Thrips.	1 bag / drum	7
Corn	Cutworm, European Corn Borer, Aphids, Beetles.	1 bag / drum	7
Apples, Pear, Peach, Apricot, Cherries, Almonds and Citrus	Green and Black Aphids, Leafminer, Stem Borer, Psylla, Leafroller, Thrips, Leafhopper.	1 bag / drum	14
Grapes	Grape Berry moth, Leafhopper.	1 bag / drum	7
Ornamentals (Open Field, Green Houses)	Leafroller, Leafminers, Bugs, Leafworm, Leafhopper, Aphids, Thrips	1 bag / drum	--

Drum = 200 lt. of water.

Bag = 100 gm of **AGRINATE-90%**

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

72 hours after the last application.

REMARKS:

- Use sprayers with low volume pressure, and use for large farms and it is not recommended for use in gardens.
- Fill 1/4 or 1/2 of the spray tank with water, then add the recommended dose of **AGRINATE-90%**, stir, fill the tank and start spraying.
- Use the recommended quantities at the beginning of infestation, repeat spraying 5-7 days intervals.
- In case there is more than one dose recommended, use the lower dosage for young crops or when the injury is slight.

COMPATIBILITY:

Compatible with most common insecticides and fungicides.

*** OTHER TRADE NAMES:**

LAMENATE – 90% SP, METHOMIL – 90% SP, CONCORD – 90% SP.

- For more details about first aid & precautions please refer to first aid & precautions index.