ASCAZINE

Water Soluble Powder

The Safe And Effective Anthelmintic Against Intestinal Round Worms



COMPOSITION: Each gm contains:

Piperazine Citrate 360 mg

PROPERTIES:

- Has a wide margin of safety, it can be used for young or pregnant animals.
- Excellent efficacy against different species of intestinal nematodes (round worms) in poultry, ruminants, equines, rabbits, dogs and cats.

INDICATIONS:

Treatment and control of ascaridiosis due to intestinal nematodes (round worms) such as: Ascaridia, Capillaria, Oxyuris, Strongylides, Oesophagostomum etc...,in poultry and other domestic animals.

DOSAGE: given with drinking water

■ **Poultry:** (Treatment duration for 2 successive days). (0.8-1) gm per bird daily. (32 mg activity / kg body wt.). (8-10 sachets of 100 gm per 1000 birds daily. OR (0.8-1) kg per 1000 birds daily. (given with drinking water)

■ Sheep, Goats, Cattle, Camels, Equines:

0.8 gm per kg body wt. given as oral drench after dissolving with proper quantity of water (110 mg activity / kg body wt.).

Rabbits:

0.5-0.8 gm per kg body wt. (70-105 mg activity / kg body wt.).

Dogs, Cats:

0.35-0.5 gm per kg body wt. (45-65 mg activity / kg body wt.).

Repeat treatment after 4-6 weeks for one day.

SAFETY PERIOD: For meat consumption: 2 days.

For eggs, milk consumption: 7 days

STORAGE: Keep in a dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.