

# AZOXIFEN-32.5%

FUNGICIDE  
SUSPENSION CONCENTRATE

VAPCO

Class:  
Strobilurin<sup>(1)</sup>  
Triazole<sup>(2)</sup>

Fungicide with protective, curative, eradicant, translaminar and systemic properties used to control a wide range of fungal diseases on a wide range of crops such as Vegetables, Fruit trees, ... etc.

**COMPOSITION:** Each liter contains:

AZOXYSTROBIN<sup>(1)</sup>  
DIFENOCONAZOLE<sup>(2)</sup>

20.0 % W/V (A.I)  
12.5 % W/V (A.I)

## PROPERTIES:

**AZOXIFEN-32.5%** : Fungicide with protective, curative, eradicant, translaminar and systemic Properties to control wide range of fungal diseases.

**AZOXIFEN-32.5%** : Strobilurin and triazole Co-Formulation with strong action .

**AZOXIFEN-32.5%** : Used in **IPM** control programmes.

**AZOXIFEN-32.5%** : Don't has any accumulated residues in treated crops.

## USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE ML / 20 LT WATER	SAFETY PERIOD (DAYS)
Tomato and Pepper	Early blight Leaf spots, Powdery mildew.	10 -15	3 7
Cucurbits except cucumber	Powdery mildew, Leaf spots and Anthracnose.	10 - 15	3
Cucumber	Powdery mildew, Downy mildew.	5	3
Potato	Early blight, Leaf spots.	10 - 15	14
Fruit Trees (Peach, Plum, Apricots, Cherries)	Powdery mildew, Leaf spots, Leaf holes.	10 - 15	14
Grapes	Powdery mildew, Leaf spots. Downy mildew, dieback	10 - 15 10	14
Citrus	Fruit rot, Dieback disease.	10	14
Strawberry	Leaf spots, Downy mildew.	5	3
Mango	Powdery mildew, Anthracnose.	12	10
Wheat and Barley	Powdery mildew, Rusts.	6	7
Peanuts	Brown spot, Rusts.	5	10

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE ML / 20 LT WATER	SAFETY PERIOD (DAYS)
Beans	Rusts.	5	7
Sugarcane	Powdery mildew, Cercospora spots.	5	10
Onion	Downey mildew, Purple blotch.	7	10

**CAUTION:**

- Used only on above mentioned crops.
- Avoid use on apple trees, and avoid spray drift to apple orchards.

**COMPATIBILITY:**

Compatible with most pesticides use the solution direct after mixing, it's preferable to make mixing trials before mixing.

**RE-ENTRY PERIOD:**

Do not enter field before 24 hours from last application.

• For more details about first aid & precautions please refer to first aid & precautions index.