

# BELATHRIN-2.5%

INSECTICIDE  
EMULSIFIABLE CONCENTRATE

VAPCO

Class:  
Pyrethroid

**Non-systemic pyrethroidal insecticide with contact and stomach action. Acts on the nervous system of insect, with rapid knockdown and long residual activity. Used to control sucking insects on a wide range of agricultural crops.**

**COMPOSITION:** Each liter contains:

**BETA-CYFLUTHRIN**

**2.5% W/V (A.I)**

**PROPERTIES:**

**BELATHRIN-2.5%:** Non-systemic pyrethroidal insecticide with contact and stomach action. Acts on the nervous system of insect.

**BELATHRIN-2.5%:** Gives rapid knockdown, and has good residual activity.

**BELATHRIN-2.5%:** Used to control a wide range of insect such as: Leafworm, Fruitworm, Stem Borer and Sucking Insects.

**BELATHRIN-2.5%:** Toxic to fish and bees.

**USES AND APPLICATION RATE:**

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Bean, Cabbage, Cauliflower, Pepper, Garlic, Tomato, Lettuce, Onion, Potato, Watermelon, Cucumber.	Garlic Moth, American Boll Worm, Cabbage Butter Flies, Cotton Leafworm.	10-12	7
Chickpeas	Budworms	4 – 8	14
Apple, Peach, Nectarine, Plum, Almond, Cherries	Leaf, Fruit and Bud Moth, Leafminer, Pear Psylla	10	7
Avocado	Fruit Spotting Bug	5 – 10	7
Cotton	African Bollworm, Diamond Bollworm, Stainer Bugs, Aphids, Whitefly, Leafworm, <i>Lygus</i> Spp.	15	---
Lucerne	Red Legged Earth Mite, Blue Oat Mite	4	---
Lupine	Cutworms, Native Budworm	4 – 8	14
	Red Legged Earth Mite, Blue Oat Mite	4	
Maize	American Boll Worm, Cotton Leafworm, Corn Borer.	10-20	14

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Sorghum	Sorghum Midge	6	14
	Native Budworm	12	
Cereals	Cutworm	6-12	7
Canola	Red Legged Earth Mite, Blue Oat Mite	4	14
	Budworms	4 – 8	
Oil Tea Tree	Pyrgo Beetle, Tea Tree Psyllid	4 - 8	12 weeks
Macadamia	Macadamia Nut Borer	10	7
	Fruit Spotting Bug	5 – 10	
Coffee	Leafminer	10	---
Ornamentals	Caterpillars, African Bollworm, Bud Fly	16	---

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

48 hours after the last application.

**COMPATIBILITY:**

Compatible with most common insecticides & fungicides.

• For more details about first aid & precautions please refer to first aid & precautions index.