

BOXER-30%

INSECTICIDE
EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Pyrethroid

Effective insecticide with contact and stomach action against many insect pests on a broad range of crops.

COMPOSITION: Each liter contains:

ETOFENPROX

30% W/V (A.I)

PROPERTIES:

BOXER-30%: Has high insecticidal activity and provides rapid killing action against various insect pests including Lepidoptera, Hemiptera, Coleoptera and Orthoptera.

BOXER-30%: Has a slight effect on various natural enemies used in Integrated Pest Management (IPM).

BOXER-30%: Has low toxicity to mammals.

BOXER-30%: Effective insecticide when used against insect pest which became resistant to Organophosphate and Carbamate compound.

BOXER-30%: Chemically stable compound and maintains its high insecticidal activity even when used in conjunction with extremely alkaline or acidic products.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Apple, Pear, Quince, Plum, Peach, Almond, Apricots, Cherries, Nectarine, Lemon, Orange, Pummelo, Mandarin, Grapefruit, Grapes, Olive trees.	Codling Moth, Grape Berry Moth, Apple Leafminers.	6.6-13.2	14
	Citrus Leafminers, Leaf Blister Moth, Smaller Tea Tortrix, Psylla, Bugs, Aphids	5-10	
Mango	Hopper	1.6-3.3	---
Cabbage, cucumber, eggplant, tomatoes	Diamond-Back Moth, Cabbage Leafworm, Cutworm, Green Peach Aphid, Aphids.	2.5-5.0	3 1
Potato	Aphids, Cutworm, Colorado Potato Beetle.	3.3-6.6	7
Tobacco	Aphids, Tobacco Cutworm, Colorado Beetle.	3.3-6.6	---

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Maize	Stem Borer, Pink Borer, Cutworm, Aphids.	3.3-6.6	7
Cotton	Bollworm, American Bollworm, Tobacco Bud Worm, Pink Bollworm, Cotton Leafworm, Cabbage Looper, Southern Armyworm, Cotton Leafroller, Aphids, Lygus Bug, Whitefly, Boll Weevil	6.6-13.3	---
Soybean	Aphids, Soybean Pod Borer, Flax Bud Worm, Bean Bug, Velvetbean Caterpillar, Bean Beetles.	2.5-5.0	14
Paddy Rice	Green Rice Leafhopper, Brown Planthopper, Rice Plant Skipper, Rice Leafminer, Rice Stem Maggot, Rice Leaf Beetle,	3.3-6.6	21
Ornamentals	Leaf Beetle, Needle Eating Beetle	4-8	---

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most fungicides and insecticides.

• For more details about first aid & precautions please refer to first aid & precautions index.