

BROMEXAN

Oral Solution
For Poultry and Animals



COMPOSITION: Each ml contains:
Bromhexine HCl **10 mg**

PROPERTIES:

- Bromhexine is an efficient mucolytic expectorant for the treatment and control of the respiratory tract disorders associated with excessive mucous secretions; as it makes them less viscous, thinner and easily removed from the respiratory passages.
- A complementary to antibiotic therapy through improvement of body response, and enhancement of their penetration into bronchial secretions; and thus considered as an adjuvant in treatment of respiratory infections.

INDICATIONS:

- **Poultry:**
As an aid in treatment of respiratory tract infections (CRD, Coryza, ILT, IB, Air succulitis), associated with excessive mucous secretions.
- **Animals:**
 - Control of coughing in cases of different pulmonary infections (Laryngitis, tracheitis, bronchitis, broncho-pneumonia, primary alveolar emphysema etc..).
 - Cases of amniotic fluid aspiration in the newly-born animals, to facilitate and expedite the expectoration of the aspired fluid.

DOSAGE:

- **Poultry:** 50-100 ml / 200 litres of drinking water for 3-5 days.
- **Bovines:** 1-2.5 ml / 50 kg B.wt. daily for 5-7 days. (0.2 – 0.5 mg activity /kg b. wt)
- **Equines:** 0.5-1.25 ml / 50 kg B.wt. daily for 7 days. (0.1 – 0.25 mg activity /kg b. wt)
- **Dogs:** 0.8-1.25 ml / 5 kg B.wt. twice daily for 5-7 days. (1.6 - 2.5 mg activity / kg b. wt)
- **Cats:** 0.5 ml / 5 kg B.wt. daily for 5-7 days. (1 mg activity /kg b. wt).

SAFETY PERIOD: for all animal products consumption: 8 hours.

STORAGE: Keep in a dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) ML.